

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
13 October 2005 (13.10.2005)

PCT

(10) International Publication Number  
**WO 2005/095383 A1**

(51) International Patent Classification<sup>7</sup>: C07D 401/14,  
A61K 31/444, A61P 25/06

TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA,  
ZM, ZW.

(21) International Application Number:  
PCT/US2005/010330

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,  
FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,  
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,  
GQ, GW, ML, MR, NE, SN, TD, TG).

(22) International Filing Date: 28 March 2005 (28.03.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/557,408 29 March 2004 (29.03.2004) US

#### Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted  
a patent (Rule 4.17(ii)) for the following designations AE,  
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,  
CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE,  
EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS,  
JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,  
MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM,  
PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY,  
TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA,  
ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ,  
NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM,  
AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT,  
BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,  
HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK,  
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG)

(71) Applicant (for all designated States except US): BRIS-  
TOL-MYERS SQUIBB COMPANY [US/US]; P.O. Box  
4000, Route 206 and Provinceline Road, Princeton, NJ  
08543-4000 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): DEGNAN, Andrew  
P. [US/US]; 123 York Street, Apt. 22P, New Haven, CT  
06511 (US). HAN, Xiaojun [CN/US]; 65 Ridgecrest  
Drive, Cheshire, CT 06410 (US). DUBOWCHIK, Gene  
M. [US/US]; 65 Spring Street, Middlefield, CT 06455  
(US). MACOR, John E. [US/US]; 20 Landon's Way,  
Guilford, CT 06437 (US).

(74) Agents: MAKUJINA, Shah et al.; Bristol-Myers Squibb  
Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).


— as to the applicant's entitlement to claim the priority of the  
earlier application (Rule 4.17(iii)) for all designations

#### Published:

— with international search report  
— before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: HETEROCYCLIC CGRP ANTAGONISTS FOR THE TREATMENT OF MIGRAINE



(57) Abstract: The present  
invention relates to compounds  
of Formula (I) as antagonists of  
calcitonin gene-related peptide  
receptors ("CGRP-receptor"),  
pharmaceutical compositions  
comprising them, methods for  
identifying them, methods of  
treatment using them and their  
use in therapy for treatment of

neurogenic vasodilation, neurogenic inflammation, migraine and other headaches, thermal injury, circulatory shock, flushing associated with menopause, airway inflammatory diseases, such as asthma and chronic obstructive pulmonary disease (COPD), and other conditions the treatment of which can be effected by the antagonism of CGRP-receptors.

## HETEROCYCLIC CGRP ANTAGONISTS FOR THE TREATMENT OF MIGRAINE

5

## FIELD OF THE INVENTION

10       The present invention relates to novel small molecule antagonists of calcitonin gene-related peptide receptors ("CGRP-receptor"), pharmaceutical compositions comprising them, methods for identifying them, methods of treatment using them and their use in therapy for treatment of neurogenic vasodilation, neurogenic inflammation, migraine, cluster headache and other  
15       headaches, thermal injury, circulatory shock, flushing associated with menopause, airway inflammatory diseases, such as asthma and chronic obstructive pulmonary disease (COPD), and other conditions the treatment of which can be effected by the antagonism of CGRP-receptors.

20

## BACKGROUND OF THE INVENTION

      Calcitonin gene-related peptide (CGRP) is a naturally occurring 37-amino-acid peptide first identified in 1982 (Amara, S. G. et al, *Science* 1982, 298, 240-244). Two forms of the peptide are expressed ( $\alpha$ CGRP and  $\beta$ CGRP) which differ by one and three amino acids in rats and humans, respectively. The peptide  
25       is widely distributed in both the peripheral (PNS) and central nervous system (CNS), principally localized in sensory afferent and central neurons, and displays a number of biological effects, including vasodilation.

      When released from the cell, CGRP binds to specific cell surface G protein-coupled receptors and exerts its biological action predominantly by  
30       activation of intracellular adenylate cyclase (Poyner, D. R. et al, *Br J Pharmacol* 1992, 105, 441-7; Van Valen, F. et al, *Neurosci Lett* 1990, 119, 195-8.). Two

- 2 -

classes of CGRP receptors, CGRP<sub>1</sub> and CGRP<sub>2</sub>, have been proposed based on the antagonist properties of the peptide fragment CGRP(8-37) and the ability of linear analogues of CGRP to activate CGRP<sub>2</sub> receptors (Juaneda, C. et al. *TiPS* 2000, 21, 432-438). However, there is lack of molecular evidence for the CGRP<sub>2</sub> receptor (Brain, S. D. et al, *TiPS* 2002, 23, 51-53). The CGRP<sub>1</sub> receptor has three components: (i) a 7 transmembrane calcitonin receptor-like receptor (CRLR); (ii) the single transmembrane receptor activity modifying protein type one (RAMP1); and (iii) the intracellular receptor component protein (RCP) (Evans B. N. et al., *J Biol Chem.* 2000, 275, 31438-43). RAMP1 is required for transport of CRLR to the plasma membrane and for ligand binding to the CGRP-receptor (McLatchie, L. M. et al, *Nature* 1998, 393, 333-339). RCP is required for signal transduction (Evans B. N. et al., *J Biol Chem.* 2000, 275, 31438-43). There are known species-specific differences in binding of small molecule antagonists to the CGRP-receptor with typically greater affinity seen for antagonism of the human receptor than for other species (Brain, S. D. et al, *TiPS* 2002, 23, 51-53). The amino acid sequence of RAMP1 determines the species selectivity, in particular, the amino acid residue Trp74 is responsible for the phenotype of the human receptor (Mallee et al. *J Biol Chem* 2002, 277, 14294-8).

Inhibitors at the receptor level to CGRP are postulated to be useful in pathophysiologic conditions where excessive CGRP receptor activation has occurred. Some of these include neurogenic vasodilation, neurogenic inflammation, migraine, cluster headache and other headaches, thermal injury, circulatory shock, menopausal flushing, and asthma. CGRP receptor activation has been implicated in the pathogenesis of migraine headache (Edvinsson L. *CNS Drugs* 2001;15(10):745-53; Williamson, D. J. *Microsc. Res. Tech.* 2001, 53, 167-178.; Grant, A. D. *Brit. J. Pharmacol.* 2002, 135, 356-362.). Serum levels of CGRP are elevated during migraine (Goadsby PJ, et al. *Ann Neurol* 1990;28:183-7) and treatment with anti-migraine drugs returns CGRP levels to normal coincident with alleviation of headache (Gallai V. et al. *Cephalalgia* 1995;15:384-90). Migraineurs exhibit elevated basal CGRP levels compared to controls (Ashina M, et al., *Pain.* 2000;86(1-2):133-8.2000). Intravenous CGRP infusion

produces lasting headache in migraineurs (Lassen LH, et al. *Cephalalgia*. 2002 Feb;22(1):54-61). Preclinical studies in dog and rat report that systemic CGRP blockade with the peptide antagonist CGRP(8-37) does not alter resting systemic hemodynamics nor regional blood flow (Shen, Y-T. et al, *J Pharmacol Exp Ther* 5 2001, 298, 551-8). Thus, CGRP-receptor antagonists may present a novel treatment for migraine that avoids the cardiovascular liabilities of active vasoconstriction associated with non-selective 5-HT<sub>1B/1D</sub> agonists, 'triptans' (e.g., sumatriptan).

There are various *in vivo* migraine models known in the literature (see De 10 Vries, P. et al, *Eur J Pharmacol* 1999, 375, 61-74). Some electrically stimulate the trigeminal ganglion and measure dilation of the intracranial vessels which they innervate (e.g., Williamson et al. *Cephalalgia* 1997 17:518-24). Since facial arteries are also innervated by the trigeminal nerve, other models study changes in facial blood flow induced by electrical trigeminal activation (e.g., Escott et al. 15 *Brain Res* 1995 669:93). Alternatively, other peripheral nerves (e.g., saphenous) and vascular beds (e.g., abdominal blood flow) are also studied (e.g., Escott et al. *Br J Pharmacol* 1993 110, 772-6;). All models have been shown to be blocked by pretreatment with the peptide antagonist CGPR(8-37) a peptide fragment that is absent the 1<sup>st</sup> seven residues, or by a small molecule CGRP-receptor antagonist. 20 In some instances, exogenous CGRP has been used as a stimulus. However, these models are all invasive terminal procedures, and none have shown the clinically important abortive effect of reversing an established increase in artery dilation or increased blood flow using post-treatment of a CGRP-receptor antagonist. Williamson et al. *Cephalalgia* 1997 17:518-24, and Williamson et al. *Cephalalgia*. 25 1997 17:525-31: used *inter alia* i.v. CGRP as a stimulus to increase intracranial dural artery diameter in sodium pentobarb anesthetized rats employing a terminal 'intravital' procedure that involved drilling to thin the skull and the creation of a closed cranial window to visualize dural arteries. The effect was blocked by pretreatment with i.v. CGRP(8-37). Escott et al. *Brain Res* 1995 669:93; *inter* 30 *alia* drilled into the rat skull and used brain electrodes to electrically stimulate the trigeminal ganglion and measured laser Doppler facial blood flow in a terminal

procedure in sodium pentobarb anesthetized rats involving neuromuscular blockade, tracheal intubation and artificial ventilation. The effect was blocked by pretreatment with CGRP(8-37). Escott et al. *Br J Pharmacol* 1993 110, 772-6; *inter alia* used intradermal (i.d.) CGRP as the stimulus to increase blood flow in

5 rat abdominal skin of sodium pentobarb anesthetized animals outfitted with cannulated jugular veins for anesthetic and drug delivery. The effect was blocked by pretreatment with i.v. CGRP(8-37). Chu et al. *Neurosci Lett* 2001 310, 169-72 used *inter alia* i.d. CGRP as the stimulus in rats and measured laser Doppler

10 changes in blood flow in the skin of the back in a terminal method using sodium pentobarb anesthetized and tracheal cannulated animals; and showed pretreatment blockade by continuous release of CGRP(8-37) from subcutaneously (s.c.) implanted osmotic pumps. Hall et al *Br J Pharmacol* 1995 114, 592-7 and Hall et al *Br J Pharmacol* 1999 126, 280-4 *inter alia* used topical CGRP to increase

15 hamster cheek pouch arteriole diameter, and i.d. CGRP to increase blood flow in rat dorsal skin of sodium pentobarb anesthetized animals outfitted with cannulated jugular veins for anesthetic and drug delivery. The effect was blocked by pretreatment with i.v. CGRP(8-37). Doods et al. *Br J Pharmacol*. 2000 Feb;129(3):420-3 *inter alia* drilled into the skull of the marmoset (new world

20 monkey) and used brain electrodes to produce electrical stimulation of the trigeminal ganglion and measured facial blood flow in an invasive terminal procedure involving neuromuscular blockade and artificial ventilation of sodium pentobarbital anesthetized primates. Increase in flow was blocked by pre-treatment of a small molecule CGRP antagonist. *See also* WO 03/272252

25 Isolated DNA Molecules Encoding Humanized Calcitonin Gene-Related Peptide Receptor, Related Non-Human Transgenic Animals and Assay Methods. Thus the method of the present invention procedure being *inter alia* a non-invasive survival model in primates measuring exogenous CGRP-induced changes in facial blood flow and demonstrating pre- and post-treatment effects of peptide and small molecule CGRP antagonists in spontaneously breathing isoflurane


30 anesthetized marmosets who recover from the procedure offers significant advantages.

- 5 -

A number of non-peptidic, small molecule CGRP-receptor antagonists have been recently reported. WO 97/09046 and equivalents disclose *inter alia* quinine and quinidine related compounds which are ligands, in particular antagonists, of CGRP-receptor. WO 98/09630 and WO 98/56779 and  
 5 equivalents disclose *inter alia* variously substituted, nitrobenzamide compounds as CGRP-receptor antagonists. WO 01/32649, WO 01/49676, and WO 01/32648 and equivalents disclose *inter alia* a series of 4-oxobutanamides and related cyclopropane derivatives as CGRP-receptor antagonists. WO 00/18764, WO 98/11128 and WO 00/55154 and equivalents disclose *inter alia* benzimidazoliny  
 10 piperidines as antagonists to CGRP-receptor. Unrelated to CGRP, a series of somatostatin antagonists have been disclosed in WO 99/52875 and WO 01/25228 and equivalents. See also U.S. 6,344,449, U.S. 6,313,097, U.S. 6,521,609, U.S. 6,552,043, US 20030181462, US20030191068 and WO 03/076432 and related applications. Thus, novel CGRP-receptor antagonists effective for the treatment  
 15 of neurogenic inflammation, migraine and other disorders would be greatly advantageous.

### SUMMARY OF THE INVENTION

Thus according to a first embodiment of the first aspect of the present  
 20 invention are provided compounds of Formula (I)



and pharmaceutically acceptable salts and solvates thereof  
 wherein

25 V is -N(R<sup>1</sup>)(R<sup>2</sup>) or OR<sup>4</sup>;

R<sup>4</sup> is H, C<sub>1-6</sub>alkyl, C<sub>1-4</sub>haloalkyl or (C<sub>1-4</sub>alkylene)<sub>0-1</sub>R<sup>4'</sup>

R<sup>4'</sup> is C<sub>3-7</sub>cycloalkyl, phenyl, adamantyl, quinuclidyl,  
 azabicyclo[2.2.1]heptyl, azetidiny, tetrahydrofuranyl,

- 6 -

furanyl, dioxolanyl, thienyl, tetrahydrothienyl, pyrrolyl,  
pyrrolinyl, pyrrolidinyl, imidazolyl, imidazolinyl,  
imidazolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl,  
oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, oxadiazolyl,  
5 thiadiazolyl, triazolyl, pyranyl, pyridyl, pyrimidinyl,  
pyrazinyl, pyridazinyl, triazinyl, piperidinyl, piperazinyl,  
morpholino, thiomorpholino or dioxolanyl; and

R<sup>4'</sup> is optionally substituted with 1 or 2 of the same or  
different substituents selected from the group  
10 consisting of halo, cyano, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>haloalkyl,  
C<sub>1-4</sub>alkoxy, hydroxy, amino, C<sub>3-7</sub>cycloalkyl,  
C<sub>1-3</sub>alkylamino, C<sub>1-3</sub>dialkylamino,  
(C<sub>1-3</sub>alkyl)<sub>0-2</sub>ureido, phenyl and benzyl; and  
R<sup>4'</sup> optionally contains 1 or 2 carbonyls wherein the carbon  
15 atom of said carbonyl is a member of the ring  
structure of R<sup>4'</sup>;

R<sup>1</sup> and R<sup>2</sup> are each independently L<sup>1</sup>, wherein L<sup>1</sup> is selected from  
the group consisting of H, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl,  
C<sub>2-6</sub>alkynyl, -C<sub>1-6</sub>alkylene-amino(C<sub>1-3</sub>alkyl)<sub>2</sub>,  
20 C<sub>3-7</sub>cycloalkyl, phenyl, azetidiny, adamantyl,  
tetrahydrofuranyl, furanyl, dioxolanyl, thienyl,  
tetrahydrothienyl, pyrrolyl, pyrrolinyl, pyrrolidinyl,  
imidazolyl, imidazolinyl, imidazolidinyl, pyrazolyl,  
pyrazolinyl, pyrazolidinyl, oxazolyl, isoxazolyl, thiazolyl,  
isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl, pyranyl,  
25 pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl,  
piperidinyl, piperazinyl, morpholino, thiomorpholino and  
dioxolanyl; and

R<sup>1</sup> and R<sup>2</sup> are each optionally and independently  
30 substituted with 1 or 2 of the same or different  
substituents selected from the group consisting of

- 7 -

halo, cyano, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>haloalkyl, C<sub>1-4</sub>alkoxy, hydroxy, amino, C<sub>3-7</sub>cycloalkyl, C<sub>1-3</sub>alkylamino, C<sub>1-3</sub>dialkylamino, (C<sub>1-3</sub>alkyl)<sub>0-2</sub>ureido, phenyl and benzyl;

5 R<sup>1</sup> and R<sup>2</sup> optionally and independently contain 1 or 2 carbonyls wherein the carbon atom of said carbonyl is a member of the heterocycles comprising R<sup>1</sup> and R<sup>2</sup>;

10 wherein L<sup>1</sup> is optionally and independently interrupted from the nitrogen to which it is attached by L<sup>2</sup>, wherein L<sup>2</sup> is independently C<sub>1-3</sub>alkylene or C<sub>1-3</sub>alkylidene; or

R<sup>1</sup> and R<sup>2</sup> together with the nitrogen to which they are attached form X,

15 wherein X is azetidiny, pyrrolyl, pyrrolinyl, pyrrolidinyl, imidazolinyl, imidazolidinyl, pyrazolinyl, pyrazolidinyl, azepinyl, diazepinyl, piperazinyl, piperidinyl, morpholino or thiomorpholino; wherein X is optionally substituted with Y, wherein

20 Y is dioxolanyl, C<sub>1-9</sub>alkyl, C<sub>2-9</sub>alkenyl, C<sub>2-9</sub>alkynyl, C<sub>1-4</sub>alkylamino, C<sub>1-4</sub>dialkylamino, C<sub>1-4</sub>alkoxy, C<sub>3-7</sub>cycloalkyl, phenyl, azetidiny, furanyl, thienyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, 25 pyrrolidinonyl, imidazolyl, imidazolinyl, imidazolidinyl, imidazolidinonyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, azepinyl, diazepinyl, pyridyl, pyrimidinyl, dihydrobenzimidazolonyl, piperazinyl, 30 piperidinyl, morpholino, benzothiazolyl, benzisothiazolyl or thiomorpholino;



- 8 -

and wherein X and Y are

optionally interrupted with Z,

wherein Z is -NHC(O)O-,  
-NHC(O)NH-, NC(O)NH<sub>2</sub>, -  
NH-, -C<sub>1-3</sub>alkylene-,  
-C<sub>1-3</sub>alkylene-, -C<sub>1-3</sub>  
alkenylene-NHC(O)O-  
C<sub>1-3</sub>alkylene-; and

optionally and independently

substituted with 1 or 2 of the  
same or different substituents  
selected from the group  
consisting of halo, C<sub>1-4</sub>alkyl,  
amino, C<sub>1-3</sub>alkylamino,  
-C<sub>1-6</sub>alkylene-  
amino(C<sub>1-3</sub>alkyl)<sub>2</sub>,  
(C<sub>1-3</sub>alkyl)<sub>0-2</sub>ureido, phenyl  
and benzyl;

X and Y optionally and

independently contain 1 or 2  
carbonyls wherein the carbon  
atom of said carbonyl is a  
member of the heterocycles  
comprising X and Y;

provided that if X is substituted with Y, and  
if X and Y are not interrupted with  
Z, then

X and Y optionally share one carbon  
atom and together form a  
spirocyclic moiety;

Q is Q' or Q";

- 9 -

wherein

Q' is  $(S^y)_sR^3$ ; and

Q'' is  $NH(S^y)_sR^3$ ,  $NHC(O)(S^y)_sR^3$ ,  $NHC(O)O(S^y)_sR^3$ ,  
 $NHC(O)NH(S^y)_sR^3$ ,  $O(S^y)_sR^3$ ,  $(S^y)_sNHR^3$ ,  
 $(S^y)_sNHC(O)R^3$ ,  $(S^y)_sNHC(O)OR^3$ ,  
 $(S^y)_sNHC(O)NHR^3$  or  $(S^y)_sOR^3$ ;

wherein  $S^y$  is  $C_{1-3}$ alkylene or  $C_{1-3}$ alkylidene and s is  
 0 or 1;

$R^3$  is  $R^{3a}$  or  $R^{3b}$

wherein

$R^{3a}$  is

- (i) a heterocycle having two fused rings with 5 to 7 members in each of said rings, said heterocycle containing one to five of the same or different heteroatoms selected from the group consisting of O, N and S and said heterocycle optionally containing 1 or 2 carbonyls wherein the carbon atom of said carbonyl is a member of said fused rings;
  - (ii) a 4 to 6 membered heterocycle containing one to three of the same or different heteroatoms selected from the group consisting of O, N and S, optionally containing 1 to 2 carbonyls, wherein the carbon atom of said carbonyl is a member of said 4 to 6 membered heterocycle;
  - (iii)  $C_{3-7}$ cycloalkyl;
  - (iv) carbazolyl, fluorenyl, phenyl, -O-phenyl, -O- $C_{1-4}$ alkylene-phenyl, or naphthyl; or
  - (v)  $C_{1-8}$ alkyl,  $C_{2-7}$ alkenyl,  $-C(O)R^{3'}$ ,  $CHC(O)O-R^{3'}$ ,  $CH(CH_3)C(O)O-R^{3'}$ ,  $-C(O)O-R^{3'}$  or  $C_{2-7}$ alkynyl;
- and

- 10 -

wherein  $R^{3a}$  is optionally substituted with 1 to 3 of the same or different substituents selected from the group consisting of benzyl, phenyl, -O-phenyl, -O- $C_{1-3}$ alkylenepheryl, - $C_{1-3}$ alkylene-OC(O)-phenyl, cyano, amino, nitro, halo,  $C_{1-6}$ alkyl,  $C_{1-3}$ mono-bi-tri-haloalkyl,  $C_{1-3}$ mono-bi-tri-haloalkyloxy, ( $C_{1-3}$ alkyl) $_{1-2}$ amine, -OR<sup>3'</sup>, -C(O)R<sup>3'</sup>, -C(O)O-R<sup>3'</sup>, -O-C(O)R<sup>3'</sup>, -N(R<sup>3'</sup>)<sub>2</sub>, -C(O)N(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)C(O)(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)C(O)N(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)C(O)OR<sup>3'</sup>, -O-C(O)N(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)SO<sub>2</sub>R<sup>3'</sup>, -SO<sub>2</sub>N(R<sup>3'</sup>)<sub>2</sub> and -SO<sub>2</sub>R<sup>3'</sup>; R<sup>3'</sup> is H or - $C_{1-6}$ alkyl; provided that if  $R^{3a}$  is , -C(O)R<sup>3'</sup>, CHC(O)O-R<sup>3'</sup>, CH(CH<sub>3</sub>)C(O)O-R<sup>3'</sup> or -C(O)O-R<sup>3'</sup>, then said -C(O)R<sup>3'</sup>, CHC(O)O-R<sup>3'</sup>, CH(CH<sub>3</sub>)C(O)O-R<sup>3'</sup> or -C(O)O-R<sup>3'</sup> are unsubstituted;

$R^{3b}$  is  $R^{3a}$  but is not phenyl, 1-naphthyl, 2-naphthyl, 1,2,3,4-tetrahydro-1-naphthyl, 1H-indol-3-yl, 1-methyl-1H-indol-3-yl, 1-formyl-1H-indol-3-yl, 1-(1,1-dimethylethoxycarbonyl)-1H-indol-3-yl, 4-imidazolyl, 1-methyl-4-imidazolyl, 2-thienyl, 3-thienyl, thiazolyl, 1H-indazol-3-yl, 1-methyl-1H-indazol-3-yl, benzo[b]fur-3-yl, benzo[b]thien-3-yl, pyridinyl, quinolinyl or isoquinolinyl; optionally substituted in the carbon skeleton with mono-, di- or trisubstituted by fluorine, chlorine or bromine atoms or by branched or unbranched alkyl groups,  $C_{3-8}$ -cycloalkyl groups, phenylalkyl groups, alkenyl, alkoxy, phenyl, phenylalkoxy,

- 11 -

- trifluoromethyl, alkoxycarbonylalkyl, carboxyalkyl,  
 alkoxycarbonyl, carboxy, dialkylaminoalkyl,  
 dialkylaminoalkoxy, hydroxy, nitro, amino,  
 acetilamino, propionylamino, benzoyl,  
 5 benzoylamino, benzoylmethylamino,  
 methylsulphonyloxy, aminocarbonyl,  
 alkylaminocarbonyl, dialkylaminocarbonyl,  
 alkanoyl, cyano, tetrazolyl, phenyl, pyridinyl,  
 thiazolyl, furyl, trifluoromethoxy,  
 10 trifluoromethylthio, trifluoromethylsulphinyl- or  
 trifluoromethylsulphonyl groups;  
 wherein said substituents may be the same or  
 different and the above-mentioned benzoyl,  
 benzoylamino- and benzoylmethylamino  
 15 groups may in turn additionally be  
 substituted in the phenyl moiety by a  
 fluorine, chlorine or bromine atom, or by an  
 alkyl, trifluoromethyl, amino or acetilamino  
 group;
- 20 D is O, NCN or NSO<sub>2</sub>C<sub>1-3</sub>alkyl;  
 A is C, N, CH or COH;  
 m and n are independently 0, 1 or 2;  
 provided that  
 if m and n are 0, then A is not N;  
 25 if m is 2, then n is not 2; or  
 if n is 2, then m is not 2;  
 E is N, CH or C;  
 p is 0 or 1;  
 if p is 1, then G, J and E together form A<sup>x</sup> or A<sup>y</sup>;  
 30 A<sup>x</sup> is a fused heterocycle having two fused rings with 5 to  
 7 members in each of said rings, said heterocycle

- 12 -

containing one to four of the same or different  
heteroatoms selected from the group consisting of  
O, N and S; and

optionally containing 1 or 2 carbonyls wherein the  
carbon atom of said carbonyl is a member  
of said fused heterocycle;

A<sup>y</sup> is a 4 to 6 membered heterocycle containing one to  
three heteroatoms selected from the group  
consisting of O, N and S; and

optionally containing 1 to 2 carbonyls, wherein the  
carbon atom of said carbonyl is a member  
of said 4 to 6 membered heterocycle;

wherein A<sup>x</sup> and A<sup>y</sup> are optionally substituted with  
C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>haloalkyl, cyano,  
C<sub>3-7</sub>cycloalkyl, phenyl, halophenyl, halo,  
furanyl, pyrrolyl, pyrrolinyl, pyrrolidinyl,  
imidazolyl, imidazoliny, imidazolidinyl,  
pyrazolyl, pyrazolinyl, pyrazolidinyl,  
pyridyl, pyrimidinyl, piperidinyl,  
piperazinyl or morpholino; or

if p is 0 such that G and J are each attached to A, then A is C, and  
G, J and A together form a spirocyclic ring system with  
said rings of said system containing A and wherein G, J  
and A together are GJA' or GJA";

wherein

GJA' is A<sup>x</sup> or A<sup>y</sup>; and

GJA" is A<sup>x</sup> or A<sup>y</sup>;

provided that

A<sup>x</sup> is not a 1,3-diaza-fused  
heterocycle; and

A<sup>y</sup> is not a 1,3-diaza-heterocycle;

- 13 -

and further provided that

if Q is Q", then  $R^3$  is  $R^{3a}$ ; and

if Q is Q', then

$R^3$  is  $R^{3b}$ ; or

5  $R^3$  is  $R^{3a}$ , p is 0 and G, J and A together form GJA".

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q' and  $R^3$  is  $R^{3b}$ .

10 According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q',  $R^3$  is  $R^{3a}$  and p is 0 such that G, J and A together form GJA".

15 According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q' and Q' is  $(S^y)_s R^3$  and s is 0.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q' and Q' is  $(S^y)_s R^3$ ,  $S^y$  is  $C_{1-3}$ alkylene and s is 1.

20 According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q' and Q' is  $(S^y)_s R^3$ ,  $S^y$  is methylene and s is 1.

25 According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q' and Q' is  $(S^y)_s R^3$ ,  $S^y$  is  $C_{1-3}$ alkylidene and s is 1.

30 According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q'.

- 14 -

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q', Q' is  $(S^y)_s R^3$ , and s is 0.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q', Q' is  $(S^y)_s R^3$ ,  $S^y$  is  $C_{1-3}$ alkylene and s is 1.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q', Q' is  $(S^y)_s R^3$ ,  $S^y$  is  $C_{1-3}$ alkylidene and s is 1.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q".

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $NH(S^y)_s R^3$ .

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $NH(S^y)_s R^3$  and s is 0.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $NH(S^y)_s R^3$ ,  $S^y$  is  $C_{1-3}$ alkylene and s is 1.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $NH(S^y)_s R^3$ ,  $S^y$  is  $C_{1-3}$ alkylidene and s is 1.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $NHC(O)(S^y)_s R^3$ .

- 15 -

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $\text{NHC(O)(S}^y\text{)}_s\text{R}^3$  and s is 0.

5       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $\text{NHC(O)(S}^y\text{)}_s\text{R}^3$ ,  $\text{S}^y$  is  $\text{C}_{1-3}$ alkylene and s is 1.

10       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $\text{NHC(O)(S}^y\text{)}_s\text{R}^3$ ,  $\text{S}^y$  is  $\text{C}_{1-3}$ alkylidene and s is 1.

15       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $\text{NHC(O)O(S}^y\text{)}_s\text{R}^3$ .

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $\text{NHC(O)O(S}^y\text{)}_s\text{R}^3$  and s is 0.

20       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $\text{NHC(O)O(S}^y\text{)}_s\text{R}^3$ ,  $\text{S}^y$  is  $\text{C}_{1-3}$ alkylene and s is 1.

25       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $\text{NHC(O)O(S}^y\text{)}_s\text{R}^3$ ,  $\text{S}^y$  is  $\text{C}_{1-3}$ alkylidene and s is 1.

30       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $\text{NHC(O)NH(S}^y\text{)}_s\text{R}^3$ .



- 16 -

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $\text{NHC(O)NH(S}^y\text{)}_s\text{R}^3$  and s is 0.

5       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $\text{NHC(O)NH(S}^y\text{)}_s\text{R}^3$ ,  $\text{S}^y$  is  $\text{C}_{1-3}$ alkylene and s is 1.

10       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and Q" is  $\text{NHC(O)NH(S}^y\text{)}_s\text{R}^3$ ,  $\text{S}^y$  is  $\text{C}_{1-3}$ alkylidene and s is 1.

15       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein V is  $\text{OR}^4$ .

      According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein V is  $\text{OR}^4$  and  $\text{R}^4$  is  $\text{C}_{1-6}$ alkyl.

20       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein V is  $-\text{N(R}^1\text{)(R}^2\text{)}$ .

      According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  
25       V is  $-\text{N(R}^1\text{)(R}^2\text{)}$  or  $\text{OR}^4$ ;

$\text{R}^4$  is H,  $\text{C}_{1-6}$ alkyl,  $\text{C}_{1-4}$ haloalkyl,  $(\text{C}_{1-4}\text{alkylene})_{0-1}\text{R}^{4'}$

$\text{R}^{4'}$  is  $\text{C}_{3-7}$ cycloalkyl, phenyl, adamantyl, quinuclidyl,

azabicyclo[2.2.1]heptyl, azetidiny, tetrahydrofuranyl,

furanyl, dioxolanyl, thienyl, tetrahydrothienyl, pyrrolyl,

30       pyrrolinyl, pyrrolidinyl, imidazolyl, imidazoliny,

imidazolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl,

oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl, pyranyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, piperidinyl, piperazinyl, morpholino, thiomorpholino or dioxolanyl; and

5  $R^{4'}$  is optionally substituted with 1 or 2 of the same or different substituents selected from the group consisting of halo, cyano,  $C_{1-4}$ alkyl,  $C_{1-4}$ haloalkyl,  $C_{1-4}$ alkoxy, hydroxy, amino,  $C_{3-7}$ cycloalkyl,  $C_{1-3}$ alkylamino,  $C_{1-3}$ dialkylamino,  $(C_{1-3}$ alkyl) $_{0-2}$ ureido, phenyl and benzyl;

$R^{4'}$  optionally contains 1 or 2 carbonyls wherein the carbon atom of said carbonyl is a member of the ring structure of  $R^{4'}$ ;

15  $R^1$  and  $R^2$  are each independently  $L^1$ , wherein  $L^1$  is selected from the group consisting of H,  $C_{1-6}$ alkyl,  $-C_{1-6}$ alkylene-amino( $C_{1-3}$ alkyl) $_2$ ,  $C_{3-7}$ cycloalkyl, phenyl, adamantyl, azetidiny, tetrahydrofuranyl, furanyl, dioxolanyl, thienyl, tetrahydrothienyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, imidazolyl, imidazoliny, imidazolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl, pyranyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, piperidinyl, piperazinyl, morpholino, thiomorpholino and dioxolanyl; and

25  $R^1$  and  $R^2$  are each optionally and independently substituted with 1 or 2 of the same or different substituents selected from the group consisting of halo, cyano,  $C_{1-4}$ alkyl,  $C_{1-4}$ haloalkyl,  $C_{1-4}$ alkoxy, hydroxy, amino,  $C_{3-7}$ cycloalkyl,  $C_{1-3}$ alkylamino,  $C_{1-3}$ dialkylamino,  $(C_{1-3}$ alkyl) $_{0-2}$ ureido, phenyl and benzyl;

30

- 18 -

R<sup>1</sup> and R<sup>2</sup> optionally and independently contain 1 or 2  
carbonyls wherein the carbon atom of said carbonyl  
is a member of the heterocycles comprising R<sup>1</sup> and  
R<sup>2</sup>;

5 wherein L<sup>1</sup> is optionally interrupted from the nitrogen to  
which it is attached by L<sup>2</sup>, wherein L<sup>2</sup> is  
C<sub>1-3</sub>alkylene; or

R<sup>1</sup> and R<sup>2</sup> together with the nitrogen to which they are attached  
form X,

10 wherein X is azetidiny, pyrroliny, pyrrolidinyl,  
imidazolinyl, imidazolidinyl, pyrazolinyl,  
pyrazolidinyl, azepiny, diazepiny, piperaziny,  
piperidinyl, morpholino or thiomorpholino;  
wherein X is optionally substituted with Y, wherein

15 Y is dioxolanyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylamino,  
C<sub>1-4</sub>dialkylamino, C<sub>1-4</sub>alkoxy,  
C<sub>3-7</sub>cycloalkyl, phenyl, azetidiny, pyrrolyl,  
pyrroliny, pyrrolidinyl, pyrrolidinonyl,  
imidazolyl, imidazolinyl, imidazolidinyl,  
20 imidazolidinonyl, pyrazolyl, pyrazolinyl,  
pyrazolidinyl, azepiny, diazepiny, pyridyl,  
pyrimidinyl, dihydrobenzimidazolonyl,  
piperaziny, piperidinyl, morpholino,  
benzothiazolyl, benzisothiazolyl or  
25 thiomorpholino;

and wherein X and Y are

optionally interrupted with Z,

wherein Z is -NHC(O)O-,  
-NHC(O)NH-, NC(O)NH<sub>2</sub>,  
30 -NH-, -C<sub>1-3</sub>alkylene-, -

- 19 -

C<sub>1-3</sub>alkylene-NHC(O)O-  
 C<sub>1-3</sub>alkylene-; and  
 optionally and independently  
 substituted with 1 or 2 of the  
 same or different substituents  
 selected from the group  
 consisting of halo, C<sub>1-4</sub>alkyl,  
 amino, C<sub>1-3</sub>alkylamino, -  
 C<sub>1-6</sub>alkylene-  
 amino(C<sub>1-3</sub>alkyl)<sub>2</sub>,  
 (C<sub>1-3</sub>alkyl)<sub>0-2</sub>ureido, phenyl  
 and benzyl;  
 X and Y optionally and  
 independently contain 1 or 2  
 carbonyls wherein the carbon  
 atom of said carbonyl is a  
 member of the heterocycles  
 comprising X and Y;  
 provided that if X is substituted with Y, and  
 if X and Y are not interrupted with  
 Z, then  
 X and Y optionally share one carbon  
 atom and together form a  
 spirocyclic moiety.

According to another embodiment of the first aspect of the present  
 invention are provided compounds according to the first embodiment of the first  
 aspect of the present invention wherein R<sup>4</sup> is H, C<sub>1-6</sub>alkyl, C<sub>1-4</sub>haloalkyl or  
 (C<sub>1-4</sub>alkylene)<sub>0-1</sub>R<sup>4'</sup>; R<sup>4'</sup> is C<sub>3-7</sub>cycloalkyl, phenyl, adamantyl, quinuclidyl,  
 azabicyclo[2.2.1]heptyl, azetidiny, tetrahydrofuranyl, furanyl, dioxolanyl,  
 thienyl, tetrahydrothienyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, imidazolyl,  
 imidazoliny, imidazolidinyl, pyrazolyl, pyrazoliny, pyrazolidinyl, oxazolyl,

- 20 -

isoxazolyl, thiazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl, pyranyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, piperidinyl, piperazinyl, morpholino, thiomorpholino or dioxolanyl; and R<sup>4'</sup> is optionally substituted with 1 or 2 of the same or different substituents selected from the group consisting of  
 5 halo, cyano, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>haloalkyl, C<sub>1-4</sub>alkoxy, hydroxy, amino, C<sub>3-7</sub>cycloalkyl, C<sub>1-3</sub>alkylamino, C<sub>1-3</sub>dialkylamino, (C<sub>1-3</sub>alkyl)<sub>0-2</sub>ureido, phenyl and benzyl.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein R<sup>4</sup> is H, C<sub>1-6</sub>alkyl, C<sub>1-4</sub>haloalkyl or  
 10 (C<sub>1-4</sub>alkylene)<sub>0-1</sub>R<sup>4'</sup>; R<sup>4'</sup> is C<sub>3-7</sub>cycloalkyl, phenyl, adamantyl, quinuclidyl, azabicyclo[2.2.1]heptyl, azetidiny, tetrahydrofuranyl, furanyl, dioxolanyl, thienyl, tetrahydrothienyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, imidazolyl, imidazoliny, imidazolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl, pyranyl,  
 15 pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, piperidinyl, piperazinyl, morpholino, thiomorpholino or dioxolanyl.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein R<sup>4</sup> is H, C<sub>1-6</sub>alkyl or (C<sub>1-4</sub>alkylene)<sub>0-1</sub>R<sup>4'</sup>;  
 20 R<sup>4'</sup> is C<sub>3-7</sub>cycloalkyl.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein V is -N(R<sup>1</sup>)(R<sup>2</sup>) and

R<sup>1</sup> and R<sup>2</sup> are each independently L<sup>1</sup>, wherein L<sup>1</sup> is selected from  
 25 the group consisting of H, C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkylene-amino(C<sub>1-3</sub>alkyl)<sub>2</sub>, C<sub>3-7</sub>cycloalkyl, phenyl, azetidiny, adamantyl, tetrahydrofuranyl, furanyl, dioxolanyl, thienyl, tetrahydrothienyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, imidazolyl, imidazoliny, imidazolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, oxazolyl, isoxazolyl, thiazolyl,  
 30 isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl, pyranyl,

- 21 -

pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl,  
piperidinyl, piperazinyl, morpholino, thiomorpholino and  
dioxolanyl; or

R<sup>1</sup> and R<sup>2</sup> together with the nitrogen to which they are attached  
form X,

wherein X is azetidiny, pyrrolinyl, pyrrolidinyl,  
imidazoliny, imidazolidinyl, pyrazoliny,  
pyrazolidinyl, azepiny, diazepiny, piperazinyl,  
piperidinyl, morpholino or Othiomorpholino;

wherein X is substituted with Y, wherein Y is

dioxolanyl, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy,  
C<sub>3-7</sub>cycloalkyl, phenyl, azetidiny, pyrrolyl,  
pyrrolinyl, pyrrolidinyl, pyrrolidinonyl,  
imidazolyl, imidazoliny, imidazolidinyl,  
imidazolidinonyl, pyrazolyl, pyrazoliny,  
pyrazolidinyl, azepiny, diazepiny, pyridyl,  
pyrimidinyl, dihydrobenzimidazolonyl,  
piperazinyl, piperidinyl, morpholino,  
benzothiazolyl, benzisothiazolyl or  
thiomorpholino;

and wherein X and Y optionally share one  
carbon atom and together form a  
spirocyclic moiety.

According to another embodiment of the first aspect of the present  
invention are provided compounds according to the first embodiment of the first  
aspect of the present invention wherein V is -N(R<sup>1</sup>)(R<sup>2</sup>) and

R<sup>1</sup> and R<sup>2</sup> are each independently L<sup>1</sup>, wherein L<sup>1</sup> is selected from  
the group consisting of H, C<sub>1-6</sub>alkyl, or

R<sup>1</sup> and R<sup>2</sup> together with the nitrogen to which they are attached  
form X,

wherein X is piperidinyl or morpholino;

- 22 -

wherein X is substituted with Y, wherein Y is  
dioxolanyl, C<sub>1-4</sub>alkyl or piperidinyl;  
and wherein X and Y optionally share one  
carbon atom and together form a  
spirocyclic moiety.

5  
According to another embodiment of the first aspect of the present  
invention are provided compounds according to the first embodiment of the first  
aspect of the present invention wherein V is -N(R<sup>1</sup>)(R<sup>2</sup>) and wherein R<sup>1</sup> and R<sup>2</sup>  
are each independently L<sup>1</sup>, wherein L<sup>1</sup> is selected from the group consisting of H,  
10 C<sub>1-6</sub>alkyl.

According to another embodiment of the first aspect of the present  
invention are provided compounds according to the first embodiment of the first  
aspect of the present invention wherein V is -N(R<sup>1</sup>)(R<sup>2</sup>) and wherein  
R<sup>1</sup> and R<sup>2</sup> together with the nitrogen to which they are attached  
15 form X,  
wherein X is piperidinyl, piperazinyl or morpholino;  
wherein X is substituted with Y, wherein Y is  
dioxolanyl, phenyl, pyridyl, piperazinyl,  
piperidinyl or C<sub>1-4</sub>alkyl;  
20 and wherein X and Y optionally share one  
carbon atom and together form a  
spirocyclic moiety.

According to another embodiment of the first aspect of the present  
invention are provided compounds according to the first embodiment of the first  
25 aspect of the present invention wherein V is -N(R<sup>1</sup>)(R<sup>2</sup>) and wherein  
R<sup>1</sup> and R<sup>2</sup> together with the nitrogen to which they are attached  
form X,  
wherein X is piperidinyl;  
wherein X is substituted with Y, wherein Y is  
30 piperidinyl.

- 23 -

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein V is  $-N(R^1)(R^2)$  and wherein

5  $R^1$  and  $R^2$  together with the nitrogen to which they are attached form X,  
wherein X is morpholino;  
wherein X is substituted with Y, wherein Y is  $C_{1-4}$ alkyl.

10 According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein V is  $-N(R^1)(R^2)$  and wherein

$R^1$  and  $R^2$  together with the nitrogen to which they are attached form X,  
wherein X is piperidinyl;  
15 wherein X is substituted with Y, wherein Y is  $C_{1-4}$ alkyl.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein V is  $-N(R^1)(R^2)$  and wherein

20  $R^1$  and  $R^2$  together with the nitrogen to which they are attached form X,  
wherein X is piperidinyl;  
wherein X is substituted with Y, wherein Y is dioxolanyl;  
25 and wherein X and Y share one carbon atom and together form a spirocyclic moiety.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein X and Y are not interrupted with Z.

30 According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first



- 24 -

aspect of the present invention wherein X and Y are not interrupted with Z; and X and Y share one carbon atom and together form a spirocyclic moiety

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^3$  is  $R^{3a}$ .

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^3$  is  $R^{3b}$ .

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^{3a}$  is a heterocycle having two fused rings with 5 to 7 members in each of said rings, said heterocycle containing one to five of the same or different heteroatoms selected from the group consisting of O, N and S.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^{3a}$  is a heterocycle having two fused rings with 5 to 7 members in each of said rings, said heterocycle containing one to five of the same or different heteroatoms selected from the group consisting of O, N and S and said heterocycle optionally containing 1 or 2 carbonyls wherein the carbon atom of said carbonyl is a member of said fused rings.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^{3a}$  is a heterocycle having two fused rings with 5 to 7 members in each of said rings, said heterocycle containing one to five of the same or different heteroatoms selected from the group consisting of O, N and S and said heterocycle optionally containing 1 or 2 carbonyls wherein the carbon atom of said carbonyl is a member of said fused rings; wherein  $R^{3a}$  is optionally substituted with 1 to 3 of the same or different substituents selected from the group consisting of benzyl, phenyl, -O-phenyl, -O- $C_{1-3}$ alkylphenyl, - $C_{1-3}$ alkylene-OC(O)-phenyl, cyano, amino, nitro, halo,  $C_{1-3}$ mono-bi-tri-haloalkyl,

- 25 -

C<sub>1-3</sub>mono-bi-tri-haloalkyloxy, C<sub>1-6</sub>alkoxy, (C<sub>1-3</sub>alkyl)<sub>1-2</sub>amine, -OR<sup>3'</sup>, -C(O)R<sup>3'</sup>, -C(O)O-R<sup>3'</sup>, -O-C(O)R<sup>3'</sup>, -N(R<sup>3'</sup>)<sub>2</sub>, -C(O)N(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)C(O)(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)C(O)N(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)C(O)OR<sup>3'</sup>, -O-C(O)N(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)SO<sub>2</sub>R<sup>3'</sup>, -SO<sub>2</sub>N(R<sup>3'</sup>)<sub>2</sub> and -SO<sub>2</sub>R<sup>3'</sup>; R<sup>3'</sup> is H or -C<sub>1-6</sub>alkyl.

5           According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein R<sup>3a</sup> is a 4 to 6 membered heterocycle containing one to three of the same or different heteroatoms selected from the group consisting of O, N and S.

10           According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein R<sup>3a</sup> is a 4 to 6 membered heterocycle containing one to three of the same or different heteroatoms selected from the group consisting of O, N and S, optionally containing 1 to 2 carbonyls, wherein  
15           the carbon atom of said carbonyl is a member of said 4 to 6 membered heterocycle.

              According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein R<sup>3a</sup> is a 4 to 6 membered heterocycle  
20           containing one to three of the same or different heteroatoms selected from the group consisting of O, N and S, optionally containing 1 to 2 carbonyls, wherein the carbon atom of said carbonyl is a member of said 4 to 6 membered heterocycle; wherein R<sup>3a</sup> is optionally substituted with 1 to 3 of the same or different substituents selected from the group consisting of benzyl, phenyl, -O-phenyl, -O-C<sub>1-3</sub>alkylphenyl, -C<sub>1-3</sub>alkylene-OC(O)-phenyl, cyano, amino, nitro,  
25           halo, C<sub>1-3</sub>mono-bi-tri-haloalkyl, C<sub>1-3</sub>mono-bi-tri-haloalkyloxy, C<sub>1-6</sub>alkoxy, (C<sub>1-3</sub>alkyl)<sub>1-2</sub>amine, -OR<sup>3'</sup>, -C(O)R<sup>3'</sup>, -C(O)O-R<sup>3'</sup>, -O-C(O)R<sup>3'</sup>, -N(R<sup>3'</sup>)<sub>2</sub>, -C(O)N(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)C(O)(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)C(O)N(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)C(O)OR<sup>3'</sup>, -O-C(O)N(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)SO<sub>2</sub>R<sup>3'</sup>, -SO<sub>2</sub>N(R<sup>3'</sup>)<sub>2</sub> and -SO<sub>2</sub>R<sup>3'</sup>; R<sup>3'</sup> is H or -C<sub>1-6</sub>alkyl.

- 26 -

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^{3a}$  is  $C_{3-7}$ cycloalkyl.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^{3a}$  is  $C_{3-7}$ cycloalkyl; wherein  $R^{3a}$  is optionally substituted with 1 to 3 of the same or different substituents selected from the group consisting of benzyl, phenyl, -O-phenyl, -O- $C_{1-3}$ alkylphenyl, - $C_{1-3}$ alkylene-OC(O)-phenyl, cyano, amino, nitro, halo,  $C_{1-3}$ mono-bi-tri-haloalkyl,  $C_{1-3}$ mono-bi-tri-haloalkyloxy,  $C_{1-6}$ alkoxy,  $(C_{1-3}alkyl)_{1-2}$ amine,  $-OR^{3'}$ ,  $-C(O)R^{3'}$ ,  $-C(O)OR^{3'}$ ,  $-O-C(O)R^{3'}$ ,  $-N(R^{3'})_2$ ,  $-C(O)N(R^{3'})_2$ ,  $-N(R^{3'})C(O)(R^{3'})_2$ ,  $-N(R^{3'})C(O)N(R^{3'})_2$ ,  $-N(R^{3'})C(O)OR^{3'}$ ,  $-O-C(O)N(R^{3'})_2$ ,  $-N(R^{3'})SO_2R^{3'}$ ,  $-SO_2N(R^{3'})_2$  and  $-SO_2R^{3'}$ ;  $R^{3'}$  is H or  $-C_{1-6}$ alkyl.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^{3a}$  is carbazolyl, fluorenyl, phenyl, -O-phenyl, -O- $C_{1-4}$ alkylene-phenyl, or naphthyl.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^{3a}$  is carbazolyl, fluorenyl, phenyl, -O-phenyl, -O- $C_{1-4}$ alkylene-phenyl, or naphthyl; wherein  $R^{3a}$  is optionally substituted with 1 to 3 of the same or different substituents selected from the group consisting of benzyl, phenyl, -O-phenyl, -O- $C_{1-3}$ alkylphenyl, - $C_{1-3}$ alkylene-OC(O)-phenyl, cyano, amino, nitro, halo,  $C_{1-3}$ mono-bi-tri-haloalkyl,  $C_{1-3}$ mono-bi-tri-haloalkyloxy,  $C_{1-6}$ alkoxy,  $(C_{1-3}alkyl)_{1-2}$ amine,  $-OR^{3'}$ ,  $-C(O)R^{3'}$ ,  $-C(O)OR^{3'}$ ,  $-O-C(O)R^{3'}$ ,  $-N(R^{3'})_2$ ,  $-C(O)N(R^{3'})_2$ ,  $-N(R^{3'})C(O)(R^{3'})_2$ ,  $-N(R^{3'})C(O)N(R^{3'})_2$ ,  $-N(R^{3'})C(O)OR^{3'}$ ,  $-O-C(O)N(R^{3'})_2$ ,  $-N(R^{3'})SO_2R^{3'}$ ,  $-SO_2N(R^{3'})_2$  and  $-SO_2R^{3'}$ ;  $R^{3'}$  is H or  $-C_{1-6}$ alkyl.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first

aspect of the present invention wherein  $R^{3a}$  is  $C_{1-8}$ alkyl,  $C_{2-7}$ alkenyl,  $-C(O)R^{3'}$ ,  $-C(O)O-R^{3'}$  or  $C_{2-7}$ alkynyl.

- According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^{3a}$  is  $C_{1-8}$ alkyl,  $C_{2-7}$ alkenyl,  $-C(O)R^{3'}$ ,  $-C(O)O-R^{3'}$  or  $C_{2-7}$ alkynyl; wherein  $R^{3a}$  is optionally substituted with 1 to 3 of the same or different substituents selected from the group consisting of benzyl, phenyl,  $-O$ -phenyl,  $-O-C_{1-3}$ alkylphenyl,  $-C_{1-3}$ alkylene- $OC(O)$ -phenyl, cyano, amino, nitro, halo,  $C_{1-3}$ mono-bi-tri-haloalkyl,  $C_{1-3}$ mono-bi-tri-haloalkyloxy,  $C_{1-6}$ alkoxy,  $(C_{1-3}$ alkyl)<sub>1-2</sub>amine,  $-OR^{3'}$ ,  $-C(O)R^{3'}$ ,  $-C(O)O-R^{3'}$ ,  $-O-C(O)R^{3'}$ ,  $-N(R^{3'})_2$ ,  $-C(O)N(R^{3'})_2$ ,  $-N(R^{3'})C(O)(R^{3'})_2$ ,  $-N(R^{3'})C(O)N(R^{3'})_2$ ,  $-N(R^{3'})C(O)OR^{3'}$ ,  $-O-C(O)N(R^{3'})_2$ ,  $-N(R^{3'})SO_2R^{3'}$ ,  $-SO_2N(R^{3'})_2$  and  $-SO_2R^{3'}$ ;  $R^{3'}$  is H or  $-C_{1-6}$ alkyl; provided that if  $R^{3a}$  is  $-C(O)R^{3'}$ ,  $CHC(O)O-R^{3'}$ ,  $CH(CH_3)C(O)O-R^{3'}$  or  $-C(O)O-R^{3'}$ , then said  $-C(O)R^{3'}$ ,  $CHC(O)O-R^{3'}$ ,  $CH(CH_3)C(O)O-R^{3'}$  or  $-C(O)O-R^{3'}$  are unsubstituted.

- According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^3$  is  $R^{3a}$  and  $R^{3a}$  is phenyl, hydroxyphenyl, azetidiny, naphthyl,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl, dihydroquinolinonyl, hydroquinolinonyl, quinolinyl, dihydroisoquinolinonyl, hydroisoquinolinonyl, isoquinolinyl, dihydroquinazolinonyl, hydroquinazolinonyl, quinazolinyl, dihydroquinoxalinonyl, hydroquinoxalinonyl, quinoxalinyl, benzimidazolyl, indazolyl, dihydrobenzimidazolonyl, hydrobenzimidazolonyl, benzimidazoliny, dihydro-benzthiazolonyl, hydrobenzthiazolonyl, benzthiazolyl, dihydrobenzoxazolyl, benzotriazolyl, dihydrobenzothiophenonyl, hydrobenzothiophenonyl, benzothienyl, dihydrobenzofuranonyl, hydrobenzofuranonyl, benzofuranyl, benzdioxolanyl, dihydroindolonyl, hydroindolonyl, indolyl, indoliziny, isoindolyl, indoliny, indazolyl, pyrazolyl, pyrazoliny, pyrazolidiny, furanyl, thienyl, pyrrolyl, pyrroliny, pyrrolidiny, imidazolyl, imidazoliny, imidazolidiny, pyridyl, puriny, carbazolyl, pyrimidiny, piperidiny, triazolopyrimidiny,

- 28 -

tetrahydropyrazolopyridinyl, piperazinyl or morpholino; optionally substituted as provided in the first embodiment of the first aspect.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^3$  is  $R^{3a}$  and  $R^{3a}$  is phenyl, naphthyl, indazolyl, benzimidazolyl, dihydrobenzoxazolyl, benzotriazolyl, benzothienyl, benzdioxolanyl, dihydroindolonyl, indolyl, furanyl, thienyl, pyridyl, purinyl, carbazolyl, piperidinyl, triazolopyrimidinyl, tetrahydropyrazolopyridinyl; optionally substituted as provided in the first embodiment of the first aspect.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^3$  is  $R^{3a}$  and  $R^{3a}$  is dihydrobenzthiazolonyl, hydrobenzthiazolonyl, benzthiazolyl, dihydrobenzothiophenonyl, hydrobenzothiophenonyl, benzothienyl, dihydrobenzofuranonyl, hydrobenzofuranonyl, benzofuranyl, dihydroindolonyl, hydroindolonyl, indolyl, indoliziny, isoindolyl, indolinyl or indazolyl; optionally substituted as provided in the first embodiment of the first aspect.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^3$  is  $R^{3a}$  and  $R^{3a}$  is dihydrobenzoxazolyl, benzotriazolyl, indolyl, halonitrophenyl, halopyrimidine, halopurinyl,  $C_{1-3}$ alkyl-nitroaminopyrimidine, triazolopyrimidinyl, pyridyl, indazolyl, phenyl or benzdioxolanyl; optionally substituted as provided in the first embodiment of the first aspect.


According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^3$  is  $R^{3a}$  and  $R^{3a}$  is naphthyl, phenyl-O-phenyl, or thienyl; optionally substituted as provided in the first embodiment of the first aspect.

- 29 -

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^3$  is  $R^{3b}$ .


According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^3$  is  $R^{3b}$  and  $R^{3b}$  is

1H-Indol-5-yl




10

1H-Indazol-5-yl




1H-Benzotriazol-5-yl




15

1,3-Dihydro-indol-2-on-5-yl




- 30 -

3H-Benzooxazol-2-on-6-yl




1,3-Dihydro-benzoimidazol-2-on-5-yl




5

1-Methyl-1,3-dihydro-benzoimidazol-2-on-6-yl




10 3,4-Dihydro-1H-quinolin-2-on-6-yl




1,4-Dihydro-benzo[d][1,3]oxazin-2-on-6-yl

- 31 -




3,4-Dihydro-1H-quinazolin-2-on-6-yl




5

3-Methyl-3,4-dihydro-1H-quinazolin-2-on-6-yl



; or

4H-Benzo[1,4]oxazin-3-on-7-yl



10

wherein  $T^y$  is H,  $C_{1-6}$ alkyl, F, Cl, Br or nitrile.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^3$  is  $R^{3b}$  and  $R^{3b}$  is azetidiny,  $C_{1-6}$ alkyl,



- 32 -

C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, dihydroquinolinonyl, hydroquinolinonyl, dihydroisoquinolinonyl, hydroisoquinolinonyl, dihydroquinazolinonyl, hydroquinazolinonyl, quinazolinyl, dihydroquinoxalinonyl, hydroquinoxalinonyl, quinoxalinyl, benzimidazolyl, 1H-indazol-5-yl, dihydrobenzimidazolonyl, 5 hydrobenzimidazolonyl, benzimidazolyl, dihydro-benzthiazolonyl, hydrobenzthiazolonyl, benzthiazolyl, dihydrobenzothiophenonyl, hydrobenzothiophenonyl, dihydrobenzofuranonyl, hydrobenzofuranonyl, benzdioxolanyl, dihydrobenzoxazolyl, benzotriazolyl, dihydroindolonyl, hydroindolonyl, indoliziny, isoindolyl, indoliny, pyrazolyl, pyrazoliny, 10 pyrazolidiny, furanyl, pyrrolyl, pyrroliny, pyrrolidiny, imidazoliny, imidazolidiny, puriny, carbazolyl, pyrimidiny, piperidiny, piperaziny or morpholino; optionally substituted as provided in the first embodiment of the first aspect.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first 15 aspect of the present invention wherein R<sup>3</sup> is R<sup>3b</sup> and R<sup>3b</sup> is dihydrobenzimidazolonyl, hydrobenzimidazolonyl, benzimidazoliny, dihydro-benzthiazolonyl, hydrobenzthiazolonyl, benzthiazolyl, dihydrobenzothiophenonyl, hydrobenzothiophenonyl, dihydrobenzofuranonyl, 20 hydrobenzofuranonyl, 1H-indazol-5-yl, benzdioxolanyl, dihydrobenzoxazolyl, benzotriazolyl, dihydroindolonyl, hydroindolonyl, indoliziny, isoindolyl, indoliny, pyrazolyl, pyrazoliny, pyrazolidiny, furanyl, pyrrolyl, pyrroliny, pyrrolidiny, imidazoliny, imidazolidiny, puriny, carbazolyl, pyrimidiny, piperidiny, piperaziny or morpholino; optionally substituted as provided in the 25 first embodiment of the first aspect.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein R<sup>3</sup> is R<sup>3b</sup> and R<sup>3b</sup> is azetidiny, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, dihydroquinolinonyl, hydroquinolinonyl, 30 dihydroisoquinolinonyl, hydroisoquinolinonyl, dihydroquinazolinonyl, hydroquinazolinonyl, quinazolinyl, dihydroquinoxalinonyl, hydroquinoxalinonyl,

quinoxaliny, benzimidazolyl, 1H-indazol-5-yl, dihydrobenzimidazolonyl, hydrobenzimidazolonyl, benzimidazoliny, dihydro-benzthiazolonyl, hydrobenzthiazolonyl, benzthiazolyl, dihydrobenzothiophenonyl, hydrobenzothiophenonyl, dihydrobenzofuranonyl, hydrobenzofuranonyl, 5 benzdioxolany, dihydrobenzoxazolyl, benzotriazolyl, puriny, carbazolyl, pyrimidinyl, piperidinyl, piperazinyl or morpholino; optionally substituted as provided in the first embodiment of the first aspect.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^3$  is  $R^{3b}$  and  $R^{3b}$  is azetidiny,  $C_{1-6}$ alkyl, 10  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl, dihydroquinolinonyl, hydroquinolinonyl, dihydroisoquinolinonyl, hydroisoquinolinonyl, dihydroquinazolinonyl, hydroquinazolinonyl, quinazolinyl, dihydroquinoxalinonyl, hydroquinoxalinonyl, quinoxaliny, benzimidazolyl, benzdioxolany, dihydrobenzoxazolyl, benzotriazolyl, dihydroindolonyl, hydroindolonyl, 1H-indazol-5-yl, indoliziny, 15 isoindolyl, indoliny, pyrazolyl, pyrazolinyl, pyrazolidiny, furanyl, pyrrolyl, pyrroliny, pyrrolidinyl, imidazoliny, imidazolidiny, puriny, carbazolyl, pyrimidinyl, piperidinyl, piperazinyl or morpholino; optionally substituted as provided in the first embodiment of the first aspect.

20 According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^3$  is  $R^{3b}$  and  $R^{3b}$  is benzdioxolany, dihydrobenzoxazolyl, benzotriazolyl, puriny, carbazolyl; optionally substituted as provided in the first embodiment of the first aspect.

25 According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $R^3$  is  $R^{3b}$  and  $R^{3b}$  is dihydrobenzoxazolyl, benzotriazolyl, indolyl, halonitrophenyl, halopyrimidinyl, halopuriny,  $C_{1-3}$ alkyl-nitroaminopyrimidinyl, triazolopyrimidinyl, pyridyl, 1H-indazol-5-yl, phenyl or 30 benzdioxolany.

- 34 -

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q' and wherein said compounds have an absolute configuration of R.

5       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q' and wherein said compounds have an absolute configuration of S.

10       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and wherein said compounds have an absolute configuration of R.

15       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein Q is Q" and wherein said compounds have an absolute configuration of S.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein m and n are each 1.

20       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein D is O.

25       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein A is C.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein A is CH.

30       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein A is N.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein E is N.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein E is CH.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein E is C.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein said compounds exhibit as described herein a CGRP Binding  $IC_{50}$  of less than 10 nM.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein said compounds exhibit as described herein a CGRP Binding  $IC_{50}$  of less than 100 nM.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein said compounds exhibit as described herein a CGRP Binding  $IC_{50}$  of less than 1000 nM.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein p is 1; and G, J and E together form  $A^x$  or  $A^y$ .

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein p is 1; and G, J and E together form  $A^x$ .

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein p is 1; and G, J and E together form  $A^y$ .

- 36 -

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein A<sup>x</sup> is a fused heterocycle having two fused rings with 5 to 7 members in each of said rings, said heterocycle containing one to four of the same or different heteroatoms selected from the group consisting of O, N and S; and optionally containing 1 or 2 carbonyls wherein the carbon atom of said carbonyl is a member of said fused heterocycle.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein A<sup>x</sup> is a fused heterocycle having two fused rings with 5 to 7 members in each of said rings, said heterocycle containing one to four of the same or different heteroatoms selected from the group consisting of O, N and S.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein A<sup>x</sup> is a fused heterocycle having two fused rings with 5 to 7 members in each of said rings, said heterocycle containing one to four of the same or different heteroatoms selected from the group consisting of O, N and S and wherein A<sup>x</sup> is substituted with phenyl.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein A<sup>x</sup> is a fused heterocycle described herein.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein A<sup>y</sup> is a 4 to 6 membered heterocycle containing one to three heteroatoms selected from the group consisting of O, N and S; and optionally containing 1 to 2 carbonyls, wherein the carbon atom of said carbonyl is a member of said 4 to 6 membered heterocycle.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first

- 37 -

aspect of the present invention wherein A<sup>y</sup> is a 4 to 6 membered heterocycle containing one to three heteroatoms selected from the group consisting of O, N and S.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein A<sup>y</sup> is a 4 to 6 membered heterocycle containing one to three heteroatoms selected from the group consisting of O, N and S; and optionally containing 1 to 2 carbonyls, wherein the carbon atom of said carbonyl is a member of said 4 to 6 membered heterocycle; and wherein A<sup>y</sup> is substituted with phenyl.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein A<sup>y</sup> is a 4 to 6 membered heterocycle described herein.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein p is 0 such that G and J are each attached to A, then G, J and A together form a spirocyclic ring system with said rings of said system containing A and wherein G, J and A together are GJA' or GJA".

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein p is 0 such that G and J are each attached to A, then G, J and A together form a spirocyclic ring system with said rings of said system containing A and wherein G, J and A together are GJA'.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein p is 0 such that G and J are each attached to A, then G, J and A together form a spirocyclic ring system with said rings of said system containing A and wherein G, J and A together are GJA".

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first

- 38 -

aspect of the present invention wherein  $p$  is 0 such that G and J are each attached to A, then G, J and A together form a spirocyclic ring system with said rings of said system containing A and wherein G, J and A together are GJA' and GJA' is  $A^x$ .

5       According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $p$  is 0 such that G and J are each attached to A, then G, J and A together form a spirocyclic ring system with said rings of said system containing A and wherein G, J and A together are GJA' and GJA' is  
10    $A^y$ .

      According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $p$  is 0 such that G and J are each attached to A, then G, J and A together form a spirocyclic ring system with said rings of  
15   said system containing A and wherein G, J and A together are GJA" and GJA" is  $A^x$ .

      According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein  $p$  is 0 such that G and J are each attached  
20   to A, then G, J and A together form a spirocyclic ring system with said rings of said system containing A and wherein G, J and A together are GJA" and GJA" is  $A^y$ .

      According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first  
25   aspect of the present invention wherein  $p$  is 0 such that G and J are each attached to A, then G, J and A together form a spirocyclic ring system with said rings of said system containing A and wherein G, J and A together are form a heterocycle selected from the group consisting of imidazolinonyl, imidazolidinonyl, dihydroquinolinonyl, dihydroisoquinolinonyl, dihydroquinazolinonyl,  
30   dihydroquinoxalinonyl, dihydrobenzoxazinyl, hydrobenzoxazinyl, dihydrobenzoxazinonyl, dihydrobenzimidazolonyl, dihydrobenzimidazolyl,

dihydro-benzthiazolonyl, dihydrobenzthiazolyl, dihydrobenzothiophenonyl, dihydrobenzofuranonyl, dihydroindolonyl, indoliny, pyrazoliny, pyrazolidiny, pyrroliny, pyrrolidiny, imidazoliny, imidazolidiny, piperidiny, piperaziny and morpholino; wherein said heterocycle is optionally substituted with C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>haloalkyl, cyano, C<sub>3-7</sub>cycloalkyl, phenyl, halophenyl, furanyl, pyrrolyl, pyrroliny, pyrrolidiny, imidazolyl, imidazoliny, imidazolidiny, pyrazolyl, pyrazoliny, pyrazolidiny, pyridyl, pyrimidiny, piperidiny, piperaziny or morpholino.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein p is 0 such that G and J are each attached to A, then G, J and A together form a spirocyclic ring system with said rings of said system containing A and wherein G, J and A together are form a heterocycle selected from the group consisting of imidazoliny, imidazolidiny, dihydroquinolinonyl, dihydroisoquinolinonyl, dihydroquinazolinonyl, dihydroquinoxalinonyl, dihydrobenzoxaziny, hydrobenzoxaziny, dihydrobenzoxazinonyl, dihydrobenzimidazolonyl, dihydrobenzimidazolyl, dihydro-benzthiazolonyl, dihydrobenzthiazolyl, dihydrobenzothiophenonyl, dihydrobenzofuranonyl, dihydroindolonyl, indoliny, pyrazoliny, pyrazolidiny, pyrroliny, pyrrolidiny, imidazoliny, imidazolidiny, piperidiny, piperaziny and morpholino; wherein said heterocycle is optionally substituted with C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>haloalkyl, cyano, C<sub>3-7</sub>cycloalkyl, phenyl, halophenyl, furanyl, pyrazolyl, pyrazoliny, pyrazolidiny, pyridyl, pyrimidiny, piperidiny, piperaziny or morpholino.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein p is 0 such that G and J are each attached to A, then G, J and A together form a spirocyclic ring system with said rings of said system containing A and wherein G, J and A together are form a heterocycle selected from the group consisting of imidazoliny, imidazolidiny, dihydroquinolinonyl, dihydroisoquinolinonyl, dihydroquinazolinonyl,



- 40 -

5 dihydrobenzofuranonyl, dihydroindolonyl, indoliny, pyrazoliny, pyrazolidiny, pyrroliny, pyrrolidiny, imidazoliny, imidazolidiny, piperidiny, piperaziny and morpholino; wherein said heterocycle is optionally substituted with C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>haloalkyl, cyano, C<sub>3-7</sub>cycloalkyl, phenyl, halophenyl, piperaziny or morpholino.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein p is 0 such that G and J are each attached to A, then G, J and A together form a spirocyclic ring system with said rings of  
10 said system containing A and wherein G, J and A together are form a heterocycle selected from the group consisting of imidazolinonyl, imidazolidinonyl, dihydroquinolinonyl, dihydroisoquinolinonyl, dihydroquinazolinonyl, dihydroquinoxalinonyl, dihydrobenzoxazinyl, hydrobenzoxazinyl, dihydrobenzoxazinonyl, dihydrobenzimidazolonyl, dihydrobenzimidazolyl,  
15 dihydro-benzthiazolonyl, dihydrobenzthiazolyl, dihydrobenzothiophenonyl, dihydrobenzofuranonyl, dihydroindolonyl, indoliny, pyrazoliny, pyrazolidiny, pyrroliny, pyrrolidiny, imidazoliny, imidazolidiny, piperidiny, piperaziny and morpholino.

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein p is 0 such that G and J are each attached to A, then G, J and A together form a spirocyclic ring system with said rings of  
20 said system containing A and wherein G, J and A together are form a heterocycle selected from the group consisting of imidazolinonyl, imidazolidinonyl, dihydroquinolinonyl, dihydroisoquinolinonyl, dihydroquinazolinonyl, dihydroquinoxalinonyl, dihydrobenzoxazinyl, hydrobenzoxazinyl and dihydrobenzoxazinonyl.  
25

According to another embodiment of the first aspect of the present invention are provided compounds according to the first embodiment of the first aspect of the present invention wherein p is 0 such that G and J are each attached to A, then G, J and A together form a spirocyclic ring system with said rings of  
30

said system containing A and wherein G, J and A together are form a heterocycle selected from the group consisting of imidazolinonyl, imidazolidinonyl, dihydroquinolinonyl, dihydroisoquinolinonyl, dihydroquinazolinonyl, dihydroquinoxalinonyl and dihydrobenzoxazinyl.

5           According to various embodiments of a second aspect of the present invention are provided pharmaceutical compositions comprising compounds of Formula (I) as defined herein.

          According to various embodiments of a third aspect of the present invention are provided methods of treating inflammation (particularly neurogenic  
10 inflammation), headache (particularly migraine), pain, thermal injury, circulatory shock, diabetes, Reynaud's syndrome, peripheral arterial insufficiency, subarachnoid/ cranial hemorrhage, tumor growth, flushing associated with menopause and other conditions the treatment of which can be effected by the antagonism of the CGRP receptor by the administration of pharmaceutical  
15 compositions comprising compounds of Formula (I) as defined herein.

          According to various embodiments of a fourth aspect of the present invention are uses of the compounds of the present invention selected from the group consisting of (a) immune regulation in gut mucosa (b) protective effect against cardiac anaphylactic injury (c) stimulating or preventing interleukin-  
20 1b(IL-1b)-stimulation of bone resorption (d) modulating expression of NK1 receptors in spinal neurons and (e) airway inflammatory diseases and chronic obstructive pulmonary disease including asthma. *See (a) Calcitonin Receptor-Like Receptor Is Expressed on Gastrointestinal Immune Cells. Hagner, Stefanie; Knauer, Jens; Haberberger, Rainer; Goeke, Burkhard; Voigt, Karlheinz; McGregor, Gerard Patrick. Institute of Physiology, Philipps University, Marburg, Germany. Digestion (2002), 66(4), 197-203; (b) Protective effects of calcitonin gene-related peptide-mediated evodiamine on guinea-pig cardiac anaphylaxis. Rang, Wei-Qing; Du, Yan-Hua; Hu, Chang-Ping; Ye, Feng; Tan, Gui-Shan; Deng, Han-Wu; Li, Yuan-Jian. School of Pharmaceutical Sciences, Department of Pharmacology, Central South University, Xiang-Ya Road 88, Changsha, Hunan, Naunyn-Schmiedeberg's Archives of Pharmacology (2003),*

25  
30

- 42 -

- 367(3), 306-311; (c) The experimental study on the effect calcitonin gene-related peptide on bone resorption mediated by interleukin-1. Lian, Kai; Du, Jingyuan; Rao, Zhenyu; Luo, Huaican. Department of Orthopedics, Xiehe Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Peop. Rep. China. Journal of Tongji Medical University (2001), 21(4), 304-307, 5 (d) Calcitonin gene-related Peptide regulates expression of neurokinin1 receptors by rat spinal neurons. Seybold VS, McCarron KE, Mermelstein PG, Groth RD, Abrahams LG. J. Neurosci. 2003 23 (5): 1816-1824. Department of Neuroscience, University of Minnesota, Minneapolis, Minnesota 55455, and 10 Department of Pharmacology, Toxicology, and Therapeutics, University of Kansas Medical Center, Kansas City, Kansas 66160 (e) Attenuation of antigen-induced airway hyperresponsiveness in CGRP-deficient mice. Aoki-Nagase, Tomoko; Nagase, Takahide; Oh-Hashi, Yoshio; Shindo, Takayuki; Kurihara, Yukiko; Yamaguchi, Yasuhiro; Yamamoto, Hiroshi; Tomita, Tetsuji; Ohga, 15 Eijiro; Nagai, Ryozi; Kurihara, Hiroki; Ouchi, Yasuyoshi. Department of Geriatric Medicine, Graduate School of Medicine, University of Tokyo, Tokyo, Japan. American Journal of Physiology (2002), 283(5,Pt. 1), L963-L970; (f) Calcitonin gene-related peptide as inflammatory mediator. Springer, Jochen; Geppetti, Pierangelo; Fischer, Axel; Groneberg, David A. Charite Campus- 20 Virchow, Department of Pediatric Pneumology and Immunology, Division of Allergy Research, Humboldt-University Berlin, Berlin, Germany. Pulmonary Pharmacology & Therapeutics (2003), 16(3), 121-130; and (g) Pharmacological targets for the inhibition of neurogenic inflammation. Helyes, Zsuzsanna; Pinter, Erika; Nemeth, Jozsef; Szolcsanyi, Janos. Department of Pharmacology and 25 Pharmacotherapy, Faculty of Medicine, University of Pecs, Pecs, Hung. Current Medicinal Chemistry: Anti-Inflammatory & Anti-Allergy Agents (2003), 2(2), 191-218 all incorporated by reference herein.

According to various embodiments of a fifth aspect of the present invention are provided combinations of the compounds of the present invention 30 with one or more agents selected from the group consisting of COX-2 inhibitors,

- 43 -

NSAIDS, aspirin, acetaminophen, triptans, ergotamine and caffeine for the treatment of migraine.

Other embodiments of the present invention may comprise a suitable combination of two or more of the embodiments and/or aspects disclosed herein.

5 Yet other embodiments of the present invention may comprise a suitable subset of an embodiment and/or aspect disclosed herein or combinations thereof.

Still yet other embodiments and aspects of the invention will be apparent according to the description provided below.

10

### DETAILED DESCRIPTION OF THE INVENTION

The description of the invention herein should be construed in congruity with the laws and principals of chemical bonding. For example, it may be necessary to remove a hydrogen atom in order accommodate a substituent at any given location.

15

As used herein, "heterocyclic" or "heterocycle" includes cyclic moieties containing one or more heteroatoms, (*e.g.*, O, N or S) said heterocycles include those that are aromatic and those that are not, *i.e.*, "alicyclic", unless otherwise specified.

20

As used herein, the term "fused bicyclic system" when describing for example a 5.6-fused bicyclic system containing 1 to 4 nitrogen atoms includes aromatic and alicyclic systems, *e.g.* indolizine, indole, isoindole, 3H-indole, indoline, indazole or benzimidazole.

25

If a substituent is named generically, then any and all species of that genus comprise that aspect of the invention. For example, a substituent generically named as "pyrrolonyl" (the radical of "pyrrolone", a pyrrole having a carbonyl) includes pyrrol-2-onyls wherein the carbonyl is adjacent to the nitrogen and pyrrol-3-onyls wherein the carbonyl and nitrogen have an intervening methylene.

30

Similarly, the present invention comprises that a substituent may be attached at any and all suitable points of attachment on said substituent unless otherwise specified.

- 44 -


However, it is also understood that the compounds encompassed by the present invention are those that are chemically stable, i.e., heteroalicyclic substituents of the present invention should not be attached in such a way that a heteroatom in said heteroalicyclic substituent is alpha to a point of attachment wherein said point of attachment is also a heteroatom.

An embodiment or aspect which depends from another embodiment or aspect, will describe only the variables having values or provisos that differ from the embodiment or aspect from which it depends. If for example a dependent embodiment only addresses  $R^2$ , then the variables and provisos not related to  $R^2$  should reflect that of the embodiment from which it depends.

If a variable is quantified with a value of zero, then a bond attaching said variable should no longer be represented.

As used herein, "alkylene" means a divalent alkane, i.e., an alkane having two hydrogen atoms removed from said alkane (said hydrogen removed from two different carbon atoms when said alkane contains more than one carbon atom), e.g.,  $-\text{CH}_2\text{CH}_2\text{CH}_2-$ .

As used herein, "alkylidene" means an alkane having two hydrogen atoms removed from one carbon atom in said alkane, e.g.,



It should be understood that the alternating double bond designations in the six-membered ring of the 5,6-membered fused structure represented in Formula (I) are relative and represent the delocalized  $\pi$  orbital electrons of said ring.

As used herein, "aryl" or "ar-" includes phenyl or naphthyl.

As used herein, "heterocyclic" or "heterocyclo" includes both heteroaryl and heteroalicyclic.

As used herein, "halo" or "halogen" includes fluoro, chloro, bromo and iodo and further means one or more of the same or different halogens may be substituted on a respective moiety.

- 45 -

Unless specified otherwise, acyclic hydrocarbons such as alkyl, alkoxy, alkenyl and alkynyl may be branched or straight chained.

It is to be understood that the present invention may include any and all possible stereoisomers, geometric isomers, diastereoisomers, enantiomers,  
5 anomers and optical isomers, unless a particular description specifies otherwise.

As used herein, "Trp74", means that the 74<sup>th</sup> residue in RAMP1 is tryptophan (Mallee et al. *J Biol Chem* 2002, 277, 14294-8) incorporated by reference herein.

As used herein "anti-migraine compound" includes any compound,  
10 peptide or peptide fragment (modified or unmodified) capable of reversing or attenuating CGRP-receptor mediated vasodilation, (e.g., CGRP-receptor antagonists).

As used herein "test compound" includes any compound, peptide or peptide fragment (modified or unmodified) being tested to determine if it is  
15 capable of reversing or attenuating CGRP-receptor mediated vasodilation, (e.g., putative CGRP-receptor antagonists).

As used herein, "CGRP-receptor agonist" includes any compound, peptide or peptide fragment (modified or unmodified) capable of inducing CGRP-receptor mediated vasodilation particularly by example  $\alpha$ CGRP or  $\beta$ CGRP; other  
20 members of the calcitonin family, e.g. adrenomedullin; N-terminal CGRP fragments, e.g. CGRP(1-12) CGRP(1-15) and CGRP(1-22); C-terminal amide (NH<sub>2</sub>) versions of CGRP e.g., CGRP(1-8+NH<sub>2</sub>), CGRP(1-13+NH<sub>2</sub>) or CGRP(1-14+NH<sub>2</sub>); and non-naturally occurring CGRP analogues e.g.,

[Ala<sup>1</sup> $\psi$ (CH<sub>2</sub>NH)Cys<sup>2</sup>]hCGRP which contains a pseudopeptide bond between  
25 Ala<sup>1</sup> and Cys<sup>2</sup>. See Maggi CA, Rovero P, Giuliani S, Evangelista S, Regoli D, Meli A. Biological activity of N-terminal fragments of calcitonin gene-related peptide. *Eur J Pharmacol.* 1990 Apr 10;179(1-2):217-9; Qing X, Wimalawansa SJ, Keith IM. Specific N-terminal CGRP fragments mitigate chronic hypoxic pulmonary hypertension in rats. *Regul Pept.* 2003 Jan 31;110(2):93-9; and  
30 Dennis T, Fournier A, St Pierre S, Quirion R. Structure-activity profile of calcitonin gene-related peptide in peripheral and brain tissues. Evidence for

- 46 -

receptor multiplicity. J Pharmacol Exp Ther. 1989 Nov;251(2):718-25  
incorporated by reference herein.

The compounds of this invention may exist in the form of  
pharmaceutically acceptable salts. Such salts may include addition salts with  
5 inorganic acids such as, for example, hydrochloric acid and sulfuric acid, and  
with organic acids such as, for example, acetic acid, citric acid, methanesulfonic  
acid, toluenesulfonic acid, tartaric acid and maleic acid. Further, in case the  
compounds of this invention contain an acidic group, the acidic group may exist  
in the form of alkali metal salts such as, for example, a potassium salt and a  
10 sodium salt; alkaline earth metal salts such as, for example, a magnesium salt and  
a calcium salt; and salts with organic bases such as a triethylammonium salt and  
an arginine salt. In the case of a sublingual formulation a saccharin salt or  
maleate salt may be of particular benefit. The compounds of the present invention  
may be hydrated or non-hydrated.

15 The compounds of this invention can be administered in such oral dosage  
forms as tablets, capsules (each of which includes sustained release or timed  
release formulations), pills, powders, granules, elixirs, tinctures, suspensions,  
syrops and emulsions. The compounds of this invention may also be  
administered intravenously, intraperitoneally, subcutaneously, or intramuscularly,  
20 all using dosage forms well known to those skilled in the pharmaceutical arts.  
The compounds can be administered alone, but generally will be administered  
with a pharmaceutical carrier selected upon the basis of the chosen route of  
administration and standard pharmaceutical practice. Compounds of this  
invention can also be administered in intranasal form by topical use of suitable  
25 intranasal vehicles, or by transdermal routes, using transdermal skin patches.  
When compounds of this invention are administered transdermally the dosage  
will be continuous throughout the dosage regimen.

While dosing from 0.01 mg/kg to 30 mg/kg is envisaged for compounds  
of the present invention, the dosage and dosage regimen and scheduling of a  
30 compounds of the present invention must in each case be carefully adjusted,  
utilizing sound professional judgment and considering the age, weight and

- 47 -

condition of the recipient, the route of administration and the nature and extent of the disease condition. In accordance with good clinical practice, it is preferred to administer the instant compounds at a concentration level which will produce effective beneficial effects without causing any harmful or untoward side effects.


5

### SYNTHESIS

Compounds of the present invention may be synthesized according to the general schemas provided below. Variables provided in the schema below are defined in accordance with the description of compounds of the above Formula  
 10 unless otherwise specified. The compounds of the present invention may be prepared according to Scheme 1 or Scheme 2. It may also be possible to use variations of said schemes to prepare the compounds of the present inventions, said variations known to those of ordinary skill in the art.

15

#### Scheme 1. Synthesis of Formula I Compounds




The synthesis described in Scheme 1 begins with a compound of Formula  
 20 II, which is an  $\alpha$ -hydroxycarboxylic acid with an appropriate hydroxyl protected terminus. Common hydroxy protecting groups (PG) include substituted benzyl group and trialkylsilyl group and their addition and removal are well known in the field. The carboxylic acid moiety of a Formula II compound is coupled with



- 48 -

an amine of formula  $\text{HNR}^1\text{R}^2$  using standard peptide coupling reagents to form an amide of Formula III. The hydroxy protecting group is removed resulting in a Formula IV compound. This compound is then coupled with an amine of Formula VIII (see below) in a carbamate forming reaction, generating a Formula I compound. Carbamate formation is conveniently carried using phosgene, disuccinimidyl carbonate, carbonyl diimidazole or other equivalents.


### Scheme 2. Synthesis of Formula I Compounds



10

The synthesis described by Scheme 2 begins with a compound of Formula V, which is an  $\alpha$ -hydroxycarboxylic acid with a protected carboxylate terminus. The protection is generally a methyl ester, but other protecting groups such as ethyl, t-butyl, and benzyl esters may also be used. The Formula V compound is coupled with an amine of Formula VIII (see below) in a carbamate forming reaction, as described above, to generate a Formula VI compound. The Formula VI compound is converted to a free acid compound of Formula VII which is then coupled with an amine of Formula  $\text{HNR}^1\text{R}^2$  to generate a Formula I compound.

20

**Scheme 3. Synthesis of Formula I Compounds**



5           The synthesis described by Scheme 3 begins with a compound of Formula VII from Scheme 2. The Formula VII compound is coupled with an alcohol, R<sup>4</sup>-OH. Such ester-forming reactions are well known in the art and can be carried out, for example, with carbodiimide coupling agents such as N,N-dicyclohexylcarbodiimide. In addition, it is often advantageous, especially for

10   esters of secondary and tertiary alcohols, to include additives that accelerate acylations such as 4-dimethylaminopyridine.

**Preparation of HNR<sup>1</sup>R<sup>2</sup> and Formula VIII amines**

Formula VIII and HNR<sup>1</sup>R<sup>2</sup> amines are commercially available, made by


15   literature methods or described herein.

**Preparation of Formula II and Formula V α-hydroxycarboxylic acid**

20

Formula II α-hydroxycarboxylic acid and Formula V α-hydroxycarboxylic acid ester may be made by methods known to one of ordinary skill in the art or made as described in Scheme 4.

## Scheme 4. Synthesis of Formula II and Formula V Compounds




- 5           The synthesis of compounds of Formula II and Formula V are described in Scheme 4. A compound of Formula X is deprotonated with a base such as diazabicycloundecene or tetramethylguanidine or other organic or inorganic bases well known in the art. An aldehyde of Formula IX is reacted with a glycine phosphonate of Formula X in a Wadsworth-Emmons coupling reaction affording
- 10   an olefin of Formula XI. The compound of Formula XI is converted to a ketoester of Formula XII by removal of the amino protecting group (PG) followed by hydrolysis using water either by itself or in conjunction with an acid such as hydrochloric acid, trifluoroacetic acid, or other organic or inorganic acids.

- 51 -

The  $\alpha$ -ketoester of Formula XII can be reduced to give a compound of Formula V by an appropriate reducing agent such as sodium borohydride, sodium cyanoborohydride, hydrogen in the presence of an appropriate catalyst such as palladium on carbon, or other reducing agents well known in the art. The hydroxyl of a compound of Formula V can be protected to give an appropriate hydroxyl protected terminus of Formula XIII. Common hydroxyl protecting groups (PG) include methoxymethyl ether, benzyloxymethylether, substituted benzyl groups and trialkylsilyl group and their addition and removal are well known in the field. A compound of Formula XIII can be converted to a compound of formula II by treatment with lithium chloride, lithium hydroxide, sodium hydroxide, or other organic or inorganic bases using water or other suitable solvents using methodology well known in the art.

Compounds of Formula II and Formula V can also be prepared as described below in Scheme 5.

**Scheme 5. Synthesis of Formula II and Formula V Compounds**




- 52 -

A compound of Formula XIV is deprotonated with a base such as diazabicycloundecene or tetramethylguanidine or other organic or inorganic bases well known in the art. An aldehyde of Formula IX is reacted with the lactate phosphonate of Formula X in a Wadsworth-Emmons coupling reaction affording an olefin of Formula XV. The compound of Formula XV is reduced to a lactate ester of Formula V by hydrogenation of the double bond. Reduction can either result in a racemic compound using, for example, hydrogenation over palladised charcoal, or a chiral compound using a chiral catalyst such as (-)-1,2-bis-((2R,5R)-2,5-diethylphospholano)benzene(cyclooctadiene)rhodium(I) tetrafluoroborate. The hydroxyl of a compound of Formula V can be protected to give an appropriate hydroxyl protected terminus of Formula XIII. Common hydroxyl protecting groups (PG) include methoxymethyl ether, benzyloxymethylether, substituted benzyl groups and trialkylsilyl group and their addition and removal are well known in the field. A compound of Formula XIII can be converted to a compound of formula II by treatment with lithium chloride, lithium hydroxide, sodium hydroxide, or other organic or inorganic bases using water or other suitable solvents using methodology well known in the art.

Compounds of Formula XI where  $R^3$  is an aromatic ring, can also be prepared as shown in Scheme 6.

#### Scheme 6. Synthesis of Formula XVII Compounds



Scheme 6 starts with an *N*-protected aminoacrylate of Formula XVI that can be coupled to a compound of Formula XVII comprising an aromatic ring to which is attached a leaving group (X) such as iodine or bromine in the presence

- 53 -


of a transition metal catalyst such as palladium (II) acetate in a non-reactive solvent with or without heating.

The syntheses of  $\alpha$ -hydroxy carboxylic acids of Formula II and  $\alpha$ -hydroxy carboxylic acid esters is well preceded in the literature and their  
5 syntheses should be known to anyone of ordinary skill in the art.

### INTERMEDIATES AND EXAMPLES

General.  $^1\text{H}$ -NMR and  $^{13}\text{C}$ -NMR spectra were run on a Bruker 500 or 300 MHz instrument and chemical shifts were reported in ppm ( $\delta$ ) with reference  
10 to tetramethylsilane ( $\delta = 0.0$ ). All evaporations were carried out under reduced pressure. Unless otherwise stated, LC/MS analyses were carried out on a Shimadzu instrument using a YMC C18 column (3 x 50 mm) employing a 2 min linear gradient of 0% to 100% solvent B in A in a 3 min run. For LC/MS and for Shimadzu Preparative HPLC system, Solvent A was: 10% methanol/90%  
15 water/0.1% trifluoroacetic acid, and solvent B was 90% methanol/10% water/0.1% trifluoroacetic acid with a UV detector set at 220 nm.

1-Benzyl-2',3'-dihydro-2'-oxospiro-[piperidine-4,4'(1'H)-quinazoline




20

Polyphosphoric acid (113 g) was heated to 100-110°C and stirred while 1-benzyl-piperidin-4-one (9.27 ml, 50 mmol) was added. Immediately afterwards, phenyl urea (9.55 g, 70. mmol) was added in portions small enough to avoid excessive foaming. The mixture was heated at 150-160°C overnight.  
25 Water (200 mL) was then added slowly to the mixture which had been allowed to cool to 100-110°C (at lower temperatures the mixture becomes too viscous to stir). The resulting solution was neutralized with 10N NaOH to ca. pH 8, and then extracted with chloroform. The organic phase was dried over magnesium

- 54 -


sulfate and then concentrated to give the crude product which was purified by flash column chromatography on silica gel (6:4 ethyl acetate/hexanes) to give the desired product (9.0 g, 58%). Mass spec.: 308.25 (MH)<sup>+</sup>.

5 2',3'-dihydro-2'-oxospiro-[piperidine-4,4'(1'H)-quinazoline



To a solution of 1-benzyl-2',3'-dihydro-2'-oxospiro-[piperidine-4,4'(1'H)-quinazoline (1.00 g) in degassed methanol ( 50 ml ) and 6N hydrochloric acid (2.0 ml) was added 10% palladized charcoal (150 mg ). The mixture was shaken on a Parr apparatus under an atmosphere of hydrogen at 60 psi overnight. LC/MS showed incomplete reaction. More 10% palladized charcoal (200 mg) was added, and the mixture was shaken for 2 more days. At that point, all starting material was consumed. The mixture was filtered and the filtrate concentrated to give 531 mg of the desired compound (64%). Mass spec.: 218.12 (MH)<sup>+</sup>.

4-Amino-4-cyano-piperidine-1-carboxylic acid *tert*-butyl ester




20 To a well stirred solution of 4-oxo-piperidine-1-carboxylic acid *tert*-butyl ester (9.0 g, 45.3 mmol) in methanol was added ammonium chloride (2.66 g, 49.8 mmol) at room temperature and stirred for 1 h. Sodium cyanide (2.44 g, 49.8 mmol) was added and stirring was continued for additional 16 h. The reaction mixture was quenched with 5% aqueous sodium hydrogencarbonate (50 mL),  
25 diluted with water, and the methanol removed by rotary evaporation. The

- 55 -

cyanoamine was extracted with methylene chloride (3x 100 mL), dried over sodium sulfate, and the solvents evaporated to give the desired compound as an oil in 91% yield. <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): δ 3.95 - 3.90 (m, 1 H), 3.80 - 3.71 (m, 1 H), 3.42 - 3.06 (m, 2 H), 2.04 - 1.94 (m, 1 H), 1.71 - 1.50 (m, 3 H). Mass spec.: 226 (MH)<sup>+</sup>.

2-Phenyl-1,3,8-triaza-spiro[4.5]dec-1-en-4-one, hydrochloride



To a solution of 4-amino-4-cyano-piperidine-1-carboxylic acid *tert*-butyl ester (1.0 g, 4.44 mmol) in methylene chloride (30 mL) was added triethylamine (1.24 mL, 8.88 mol), followed by benzoyl chloride (936 mg, 6.66 mmol). After 30 min, 4-(dimethylamino)pyridine (40 mg, 0.33 mmol) was added and stirring continued for additional 12 h. The reaction mixture was then quenched with 1M sodium hydroxide (10 mL), diluted with ethyl acetate (100 mL), and separated. The organic layer was washed sequentially with 1M sodium hydroxide (40 mL), aqueous sodium hydrogencarbonate (50 mL), and brine (50 mL) then dried over sodium sulfate. The desired product, 4-benzoylamino-4-cyano-piperidine-1-carboxylic acid *tert*-butyl ester was obtained in 90% yield through crystallization using 30% ethyl acetate in hexane as a solvent.


To a solution of 4-benzoylamino-4-cyano-piperidine-1-carboxylic acid *tert*-butyl ester (1.3 g, 4 mmol) in ethanol (10 mL) was added 6M sodium hydroxide (1.5 mL) followed by 30% hydrogen peroxide. The reaction mixture was then refluxed for 3 h. The reaction mixture was then diluted with water (30 mL), and the ethanol removed. The residue was diluted with ethyl acetate (100 mL). The organic phase was washed with brine (30 mL) and dried over sodium sulfate. The desired product, 4-oxo-2-phenyl-1,3,8-triaza-spiro[4.5]dec-1-ene-8-



- 56 -

carboxylic acid *tert*-butyl ester was obtained in 80% yield through crystallization from 30% ethyl acetate in hexane. The *tert*-butyl ester was then dissolved in methylene chloride (5 mL) and a saturated solution of hydrogen chloride in dioxane (25 mL) was added. After 2 h, the solvent was removed to give 2-phenyl-  
5 1,3,8-triaza-spiro[4.5]dec-1-en-4-one, hydrochloride as white powder in 95% yield. <sup>1</sup>H-NMR (500 MHz, CD<sub>3</sub>OD): δ 8.23 - 8.21 (m, 2 H), 7.96 - 7.92 (m, 1 H), 7.79 - 7.76 (m, 2 H), 3.68 - 3.64 (m, 3 H), 3.31 - 3.30 (m, 1 H), 2.47 - 2.44 (m, 4 H). Mass spec.: 230 (MH)<sup>+</sup>.

10 5-Formyl-indazole-1-carboxylic acid *tert*-butyl ester




A methylene chloride (2 mL) solution of di-*tert*-butyldicarbonate (388 mg, 1.78 mmol) was added dropwise at room temperature to a solution of 1*H*-  
15 indazole-5-carbaldehyde (273 mg, 1.87 mmol), 4-dimethylaminopyridine (114 mg, 0.94 mmol), and triethylamine (0.26 mL, 1.87 mmol) in methylene chloride (10 mL). The resulting bright yellow solution was stirred at room temperature for 16 h. Solvents were removed *in vacuo* and the residue was subjected to flash chromatography with silica gel (25 g) and ethyl acetate/hexanes (1:1) containing  
20 1% triethylamine as eluent to afford the title compound as a brownish yellow liquid (414 mg, 90%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 10.08 (s, 1H), 8.38 (s, 1H), 8.34 (s, 1H), 8.25 (d, J = 8.5 Hz, 1H), 8.04 (d, J = 8.8 Hz, 1H), 1.71 (s, 9H). <sup>13</sup>CNMR (CDCl<sub>3</sub>, 125 MHz) δ 191.8, 149.0, 142.5, 140.6, 133.0, 128.3, 126.4, 125.8, 115.3, 85.7, 27.8.

25

5-(2-Benzyloxycarbonylamino-2-methoxycarbonyl-vinyl)-indazole-1-carboxylic acid *tert*-butyl ester


- 57 -



A solution of *N*-(benzyloxycarbonyl)- $\alpha$ -phosphonoglycine trimethyl ester (5.50 g, 16.6 mmol) and tetramethylguanidine (1.99 mL, 15.9 mmol) in  
 5 anhydrous tetrahydrofuran (50 mL) was stirred at  $-78^{\circ}\text{C}$  for 20 min. To this was added a solution of 5-formyl-indazole-1-carboxylic acid *tert*-butyl ester (3.72 g, 15.1 mmol) in tetrahydrofuran (25 mL) slowly via syringe over 10 min. The reaction mixture was stirred at  $-78^{\circ}\text{C}$  for 4 h and then allowed to warm to room temperature overnight. The solvent was evaporated and the resulting residue  
 10 subjected to flash column chromatography on silica gel (1:2 ethyl acetate/hexane) giving the title compound as a white foam (5.77 g, 85%).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  8.09 (d,  $J = 9.0$  Hz, 1H), 8.08 (s, 1H), 7.84 (s, 1H), 7.67 (d,  $J = 9.0$  Hz, 1H), 7.47 (s, 1H), 7.30 (br s, 5H), 6.43 (br s, 1H), 5.09 (s, 2H), 3.84 (s, 3H), 1.72 (s, 9H). Mass spec.: 452 ( $\text{MH}$ ) $^+$ .

15

#### 2-Trimethylsilylanyl-ethanesulfonyl chloride




Sulfuryl chloride (43 mL, 539 mmol) was added in 3 min to a clear  
 20 solution of triphenylphosphine (129 g, 490 mmol) in methylene chloride (200 mL) at  $0^{\circ}\text{C}$  in a flame-dried three-neck round bottom flask. After stirring at  $0^{\circ}\text{C}$  for 5 min, the ice-water bath was removed and sodium 2-trimethylsilyl-ethanesulfonate (50 g, 245 mmol) was added in portions over 10 min. The resulting white suspension was stirred at room temperature for 16 h,  
 25 then it was filtered through a pad of celite. The filtrate was concentrated to ca 50

- 58 -

mL, ethyl acetate/hexanes (1:3, 1000 mL) and celite (40 g) were added. The mixture was stirred at room temperature for 15 min and filtered through a pad of celite. Solvents were removed *in vacuo* and the residue was loaded onto a pre-wetted column with silica gel (300 mL) using 1:3 ethyl acetate/hexanes as the eluent. Solvents were removed and the title compound was obtained as a light tan liquid (41.9g, 85%). If not used immediately, the final product should be stored under nitrogen in the freezer or refrigerator to minimize decomposition. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 3.61-3.57 (m, 2H), 1.32-1.27 (m, 2H), 0.10 (s, 9H).

10 1-(2-Trimethylsilanyl-ethanesulfonyl)-1*H*-indole-5-carboxylic acid ethyl ester




A solution of 1*H*-indole-5-carboxylic acid ethyl ester (10.31g, 58.8 mmol) in dimethylformamide (50 mL) was added dropwise at 0°C to a mixture of sodium hydride (1.83g, 76.4 mmol) in dimethylformamide (150 mL). The resulting mixture was stirred at 0°C for 30 min, then a solution of 2-trimethylsilanyl-ethanesulfonyl chloride (17.7 g, 88.2 mmol) in dimethylformamide (100mL) was added slowly at 0°C to the above mixture. After 2 h, sat. aqueous ammonium chloride (200 mL) was added, and the mixture was extracted with ethyl acetate (300 mL). After separation, the aqueous layer was extracted with ethyl acetate (2 × 150 mL). The combined organic layers were washed with brine (3 × 150 mL), and dried over anhydrous sodium sulfate. Solvents were removed *in vacuo* and the residue was subjected to flash chromatography on silica gel using 1:1.5 methylene chloride/hexanes as eluent to afford the title compound as a white solid (15.8 g, 79%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 8.36 (d, *J* = 1.5 Hz, 1H), 8.03 (dd, *J* = 9.0, 2.0 Hz, 1H), 7.92 (d, *J* = 8.5 Hz, 1H), 7.50 (d, *J* = 3.5 Hz, 1H), 6.75 (d, *J* = 3.5 Hz, 1H), 3.94 (s, 3H), 3.21 –

- 59 -

3.18 (m, 2H), 0.84 – 0.80 (m, 2H), -0.06 (s, 9H).  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ , 125 MHz)  $\delta$  167.3, 137.7, 130.3, 128.3, 125.9, 125.5, 124.0, 112.8, 108.3, 52.2, 51.2, 10.1, -2.1. Mass spec. 354.12 (MH) $^+$ .


5 Similarly prepared:

1-(2-Trimethylsilyl-ethanesulfonyl)-1*H*-indazole-5-carboxylic acid ethyl ester



$^1\text{H}$ -NMR ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  8.51 (s, 1H), 8.34 (s, 1H), 8.21 (dd,  $J$  = 8.9, 1.5 Hz, 1H), 8.12 (d,  $J$  = 9.2 Hz, 1H), 3.96 (s, 3H), 3.42 – 3.39 (m, 2H), 0.86 – 0.82 (m, 2H), -0.02 (s, 9H).  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ , 125 MHz)  $\delta$  166.4, 143.1, 141.2, 130.1, 126.5, 125.0, 124.2, 112.9, 52.5, 51.3, 9.8, -2.1. Mass spec. 355.13 (MH) $^+$ .

[1-(2-Trimethylsilyl-ethanesulfonyl)-1*H*-indol-5-yl]-methanol




15

A solution of diisobutylaluminum hydride (82.9 mL, 1M in toluene, 82.9 mmol) was added slowly at 0°C to the solution of 1-(2-trimethylsilyl-ethanesulfonyl)-1*H*-indole-5-carboxylic acid ethyl ester (8.81 g, 25.9 mmol) in toluene (200mL). After it was stirred at 0°C for 45 min, the reaction was quenched by the addition of methanol (26mL), pulverized sodium sulfate decahydrate (194 g) and celite (26 mL). The mixture was warmed up to room temperature in 1h and filtered through a pad of celite. Solvents were removed *in*

- 60 -

*vacuo* to afford the title compound as a very viscous liquid, which solidified upon cooling. A white solid (8.08 g, 100% yield).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  7.87 (d,  $J = 8.5$  Hz, 1H), 7.62 (s, 1H), 7.44 (d,  $J = 3.7$  Hz, 1H), 7.35 (dd,  $J = 8.6$ , 1.5 Hz, 1H), 6.66 (d,  $J = 3.7$  Hz, 1H), 4.79 (s, 2H), 3.18 – 3.14 (m, 2H), 1.73 (s, 1H),  
 5 0.85 – 0.82 (m, 2H), -0.06 (s, 9H). Mass spec. 312.14 (MH) $^+$ .


[1-(2-Trimethylsilyl-ethanesulfonyl)-1*H*-indazol-5-yl]-methanol



10 A solution of 1-(2-trimethylsilyl-ethanesulfonyl)-1*H*-indazole-5-carboxylic acid ethyl ester (azeotropically dried with toluene (2x), 5.77g, 16.9 mmol) in tetrahydrofuran (50 mL) was added at 0°C to a mixture of lithium borohydride (3.68g, 169 mmol) in tetrahydrofuran (100 mL). The mixture was warmed up to room temperature and stirred for 14h. It was cooled to 0°C and  
 15 lithium borohydride (3.5g) was added. The mixture was warmed up to room temperature and stirred for 14h. It was re-cooled to 0°C and sat. aqueous ammonium chloride (25 mL) was added slowly. The resulted white suspension was filtered through a pad of celite, solvents were removed and the residue was subjected to flash chromatography using ethyl acetate/hexanes (1:1.5) with 1%  
 20 triethylamine to afford the title compound as a white solid (3.8g, 72%).  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , 500 MHz)  $\delta$  8.41 (s, 1H), 8.04 (d,  $J = 8.5$  Hz, 1H), 7.85 (s, 1H), 7.61 (dd,  $J = 8.5$ , 1.2 Hz, 1H), 4.76 (s, 2H), 3.49 – 3.46 (m, 2H), 0.76 – 0.72 (m, 2H), -0.03 (s, 9H);  $^{13}\text{C-NMR}$  ( $\text{CD}_3\text{OD}$ , 125 MHz)  $\delta$  141.2, 140.9, 138.3, 129.2, 125.8, 119.6, 112.7, 63.8, 50.8, 9.9, -3.2. Mass spec. 313.12 (MH) $^+$ .

25


- 61 -

1-(2-Trimethylsilyl-ethanesulfonyl)-1*H*-indole-5-carbaldehyde

A solution of [1-(2-trimethylsilyl-ethanesulfonyl)-1*H*-indol-5-yl]-  
5 methanol (2.1 g, 6.74 mmol) in methylene chloride (30 mL) was added at 0°C to a  
mixture of activated manganese dioxide (22g, azeotropically dried with toluene  
(2x)) and methylene chloride (70 mL) in a 500 mL round bottom flask. The  
reaction mixture was stirred at 0°C for 30 min and filtered through a pad of celite.  
Solvents were removed *in vacuo* to afford the title compound as a white solid  
10 (1.8g, 80%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 10.06 (s, 1H), 8.15 (s, 1H), 8.01 (d, J  
= 8.6 Hz, 1H), 7.87 (dd, J = 8.6, 1.5 Hz, 1H), 7.54 (d, J = 3.4 Hz, 1H), 6.80 (d, J =  
3.6 Hz, 1H), 3.24 – 3.20 (m, 2H), 0.86 – 0.82 (m, 2H), -0.06 (s, 9H). <sup>13</sup>C-NMR  
(CDCl<sub>3</sub>, 125 MHz) δ 191.9, 138.5, 132.3, 130.7, 128.8, 125.3, 125.1, 1134.6,  
108.4, 51.4, 10.2, -2.1. Mass spec. 310.12 (MH)<sup>+</sup>.

15


Similarly prepared:

1-(2-Trimethylsilyl-ethanesulfonyl)-1*H*-indazole-5-carbaldehydeMass spec. 311.10 (MH)<sup>+</sup>.

20

- 62 -

2-Benzoyloxycarbonylamino-3-[1-(2-trimethylsilyl-ethanesulfonyl)-1H-indol-5-yl]-acrylic acid methyl ester



5           1,1,3,3-Tetramethylguanidine (0.68 mL, 5.43 mmol) was added at room temperature to a solution of N-(benzyloxycarbonyl)- $\alpha$ -phosphonoglycine trimethyl ester (1.88g, 5.69 mmol) in tetrahydrofuran (40 mL). The mixture was stirred at room temperature for 15 min and cooled to  $-78^{\circ}\text{C}$ , and a solution of 1-(2-trimethylsilyl-ethanesulfonyl)-1H-indole-5-carbaldehyde (1.6g, 5.17 mmol) in

10 tetrahydrofuran (15 mL) was added slowly. The resulting reaction mixture was stirred at  $-78^{\circ}\text{C}$  for 2h and then warmed to room temperature in 3h. Solvents were removed *in vacuo* and the residue was subjected to flash chromatography on silica gel using methylene chloride/hexanes (1:1.5) with 1% triethylamine as eluent to afford the title compound as a 92:8 Z/E mixture (determined by


15 integration of  $\text{CO}_2\text{CH}_3$ , for Z isomer at 3.79 ppm, and E isomer at 3.65 ppm). For the Z isomer:  $^1\text{H-NMR}$  ( $\text{CD}_3\text{CN}$ , 500 MHz)  $\delta$  7.96 (s, 1H), 7.91 (d,  $J = 8.5$  Hz, 1H), 7.66 (d,  $J = 8.5$  Hz, 1H), 7.56 (d,  $J = 3.7$  Hz, 1H), 7.51 (s, 1H), 7.43 – 7.35 (m, 5H), 7.67 (d,  $J = 3.7$  Hz, 1H), 5.16 (s, 2H), 3.79 (s, 3H), 3.42 – 3.38 (m, 2H), 0.87 – 0.83 (m, 2H), -0.04 (s, 9H). Mass spec. 515.20 (MH) $^+$ .

20

Similarly prepared:

2-Benzoyloxycarbonylamino-3-[1-(2-trimethylsilyl-ethanesulfonyl)-1H-indazol-5-yl]-acrylic acid methyl ester


- 63 -



Flash chromatography on silica gel using methylene chloride containing 1% triethylamine as eluent afforded the title compound as a 95:5 Z/E mixture (determined by the integration of  $-\text{CH}=\text{C}(\text{CO}_2\text{Me})(\text{NHCBz})$ , 3.72g, 92%). For the Z isomer:  $^1\text{H-NMR}$  ( $\text{CD}_3\text{CN}$ , 500 MHz)  $\delta$  8.39 (s, 1H), 8.12 (s, 1H), 8.03 (d,  $J = 8.8$  Hz, 1H), 7.84 (dd,  $J = 8.8, 1.2$  Hz, 1H), 7.51 (s, 1H), 7.43 – 7.35 (m, 5H), 5.14 (s, 2H), 3.81 (s, 3H), 3.51 – 3.47 (m, 2H), 0.83 – 0.79 (m, 2H), -0.02 (s, 9H). Mass spec. 516.18 (MH) $^+$ .

10

7-Methyl-2-(2-trimethylsilyl-ethanesulfonyl)-2H-indazole-5-carbaldehyde




To a suspension of 7-methylindazole 5-aldehyde (3.0 g, 18.7 mmol) in methylene chloride (150 mL) was added triethylamine (7.83 mL, 56.2 mL, 3 equiv) followed by dropwise addition of neat 2-trimethylsilyl-ethanesulfonyl chloride (5.60 g, 28.1 mmol, 1.5 equiv). The mixture gradually became homogeneous and was allowed to stir at room temperature for 16 h. The solution was concentrated to a minimum amount of methylene chloride and then subjected to flash column chromatography on silica gel (1:4 ethyl acetate/hexanes) to give 4.7 g (77%) of the product as a pale yellow solid.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ , 300 MHz)  $\delta$



- 64 -

9.98 (s, 1H), 8.77 (s, 1H), 8.09 (s, 1H), 7.64 (s, 1H), 3.64-3.58 (m, 2H), 2.65 (s, 3H), 0.88-0.82 (m, 2H), 0.01 (s, 9H).

2-Benzyloxycarbonylamino-3-[7-methyl-2-(2-trimethylsilyl-ethanesulfonyl)-  
5 2*H*-indazol-5-yl]-acrylic acid methyl ester



To a solution of N-(benzyloxycarbonyl)- $\alpha$ -phosphonoglycine trimethyl ester (4.93 g, 14.9 mmol, 1.1 equiv) in anhydrous tetrahydrofuran (75 mL) was  
10 added tetramethylguanidine (1.78 mL, 1.05 equiv). The mixture was stirred at room temperature under nitrogen for 5 min and was then cooled to -78°C. After stirring for 15 min at -78°C, a solution of 7-methyl-2-(2-trimethylsilyl-ethanesulfonyl)-2*H*-indazole-5-carbaldehyde in tetrahydrofuran (25 mL) was added. The reaction mixture was allowed to slowly warm to room temperature  
15 overnight. Although the reaction was incomplete, the solvent was evaporated. The resulting residue was dissolved in ethyl acetate and washed with 1M sulfuric acid. The organic layer was separated, dried over magnesium sulfate, filtered and evaporated. Flash column chromatography (1:4 ethyl acetate/hexanes) gave 2.66 g (37 %) of the product as white glass foam. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz)  $\delta$  8.48  
20 (s, 1H), 7.62 (s, 1H), 7.38-7.25 (m, 7H), 6.48 (bs, 1H), 5.10 (s, 2H), 3.83 (s, 3H), 3.58-3.52 (m, 2H), 2.51 (s, 3H), 0.89-0.83 (m, 2H), 0.02 (s, 9H). Mass spec.: 530 (MH)<sup>+</sup>.


4-Bromo-2,6-dimethylphenyldiazo-t-butyl sulfide

- 65 -



4-Bromo-2,6-dimethylaniline (20.00 g, 100 mmol) was ground to a powder with a mortar and pestle and then suspended in 24% hydrochloric acid (41 mL). The stirred mixture was cooled to -20°C and treated with sodium nitrite (7.24 g, 1.05 equiv) in water (16 mL), dropwise over 40 min while the temperature was maintained below -5°C. After a further 30 min at -5°C to -20°C, the mixture was buffered to ca. pH 5 with solid sodium acetate. This mixture (kept at ca. -10°C) was added in portions to a stirred solution of t-butyl thiol (11.3 mL, 1 equiv) in ethanol (100 mL) at 0°C over ca. 10 min. Following addition, the mixture was stirred at 0°C for 30 min and then crushed ice (ca. 150 mL) was added. The mixture was stored in the refrigerator overnight. The resulting light-brown solid was collected by filtration, washed with water, and dried under high vacuum for several hours. (26.90 g, 89%). The compound appeared to be stable as a solid but underwent significant decomposition when recrystallization from ethanol was attempted. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 1.58 (9H, s), 1.99 (6H, s), 7.21 (2H, s). Mass spec.: 303.05 (MH)<sup>+</sup>.

#### 5-Bromo-7-methylindazole




20

Into a flame-dried round bottom flask, 4-bromo-2,6-dimethylphenyldiazo-t-butyl sulfide (12.50 g, 41.5 mmol) and potassium t-butoxide (46.56 g, 10 equiv)

- 66 -

were combined. A stir bar was added and the mixture placed under nitrogen. To this was added dry DMSO (120 mL). The mixture was stirred vigorously overnight at rt. The reaction mixture was then carefully poured into a mixture of crushed ice (400 mL) and 10% hydrochloric acid (200 mL). The resulting suspension was left to stand at 4°C overnight and the solid was collected by filtration and washed with water. The crude solid was dissolved in 5:1 methylene chloride/methanol and the solution dried over magnesium sulfate and evaporated to give the product as an off-white solid (7.60 g, 87%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>/CD<sub>3</sub>OD, 500 MHz) δ 2.51 (3H, s), 7.22 (1H, s), 7.69 (1H, s), 7.94 (1H, s). Mass spec.: 211.03 (MH)<sup>+</sup>.

## 7-methylindazole-5-carboxaldehyde




5-Bromo-7-methylindazole (6.10 g, 28.9 mmol) and sodium hydride (60% in mineral oil, 1.27 g, 1.1 equiv) were weighed into a flame-dried round-bottom flask containing a magnetic stir bar. Under a nitrogen atmosphere at room temperature, dry tetrahydrofuran (30 mL) was added. The mixture was stirred at room temperature for 15 min, during which time it became homogeneous. The stirred mixture was cooled to -70°C and a solution of *sec*-butyllithium in cyclohexane (1.4M, 45 mL, 2.2 equiv) was added over several minutes. After 1 h at -70°C, dimethylformamide (10 mL) was added over several minutes. The mixture was allowed to warm to room temperature and was stirred overnight. It was then cooled to 0°C and carefully treated with 1N hydrochloric acid (60 mL). After a few minutes, solid sodium bicarbonate was added to basify the mixture to pH 9-10. The layers were separated and the aqueous phase washed twice with ethyl acetate. The combined organic phases were extracted with 0.8M sodium hydrogen sulfate (3 x 125 mL). The combined aqueous phases were washed with

- 67 -

ethyl acetate (100 mL) and then the pH was adjusted to ca. 10 with solid sodium hydroxide. The resulting suspension was extracted with ethyl acetate (3 x 150 mL). The combined organic phases were washed with brine, dried (magnesium sulfate) and evaporated to give the product as a light-tan solid (3.01 g, 65%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 2.63 (3H, s), 7.73 (1H, s), 8.12 (1H, s), 8.25 (1H, s), 10.03 (1H, s). Mass spec.: 161.06 (MH)<sup>+</sup>.

2-Benzyloxycarbonylamino-3-(7-methyl-1H-indazol-5-yl)-acrylic acid methyl ester




10

A stirred solution of N-benzyloxycarbonyl-α-phosphonoglycine trimethyl ester (5.51 g, 1.2 equiv.) in tetrahydrofuran (30 mL) at room temperature was treated with tetramethylguanidine (1.91 mL, 1.1 equiv). After 10 min, 7-methylindazole-5-carboxaldehyde (2.22 g, 13.86 mmol) in tetrahydrofuran (20 mL) was added. Disappearance of starting material was monitored by TLC and LC/MS. After 5 days at room temperature, the solvent was evaporated and the residue dissolved in ethyl acetate. The solution was washed with 2% phosphoric acid and brine, dried (magnesium sulfate) and evaporated. The residue was purified by flash chromatography on silica gel, eluting with 1) 1:1 and 2) 2:1 ethyl acetate/hexane, to give the product as a colorless foam (4.93 g, 97%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 2.43 (3H, s), 3.80 (3H, s), 5.12 (2H, s), 6.66 (1H, s), 7.28 (5H, brs), 7.33 (1H, s), 7.47 (1H, s), 7.74 (1H, s), 7.96 (1H, s). Mass spec.: 366.16 (MH)<sup>+</sup>.

25


*tert*-Butyl (Z)-1-(methoxycarbonyl)-2-(2-((2-(trimethylsilyl)ethoxy)methyl)-7-methyl-2H-indazol-5-yl)vinylcarbamate

- 68 -



To a solution of 2-((2-(trimethylsilyl)ethoxy)methyl)-7-methyl-2H-indazole-5-carbaldehyde (4.46 g, 15.4 mmol) and N-(tert-butoxycarbonyl)-methyl-2-(dimethylphosphono)glycinate (4.80 g, 1.0 equiv) in tetrahydrofuran (40 mL) at room temperature was added N,N,N',N'-tetramethylguanidine (3.29 mL, 1.7 equiv). The reaction was allowed to stir at room temperature for 3 d. The reaction was diluted with ethyl acetate and water. Poured into diethyl ether, and washed with water (2X), then brine, dried over magnesium sulfate and concentrated. Column chromatography (30% ethyl acetate/hexanes -> 40% ethyl acetate/hexanes) gave 5.90 g (83%) as a foam. Mass spec.: 462.40 (MH)<sup>+</sup>.

Methyl 3-(2-((2-(trimethylsilyl)ethoxy)methyl)-7-methyl-2H-indazol-5-yl)-2-hydroxypropanoate



15

To a solution of tert-butyl (Z)-1-(methoxycarbonyl)-2-((2-(trimethylsilyl)ethoxy)methyl)-7-methyl-2H-indazol-5-yl)vinylcarbamate (200 mg, 0.43 mmol) in dichloromethane (2 mL) at 0°C was added trifluoroacetic acid (1 mL). The ice bath was removed. After 30 minutes, the reaction was poured


20

- 69 -

into separatory funnel containing ethyl acetate and water, neutralized with solid sodium bicarbonate, and the layers separated. The organics were washed with saturated sodium bicarbonate, then brine, dried over magnesium sulfate, and concentrated. The yellow residue was treated with sodium cyanoborohydride  
5 (200 mg, 7.4 equiv) and tetrahydrofuran (2 mL). The reaction was stirred at room temperature overnight, diluted with ethyl acetate, washed with water (2X), then brine, dried over magnesium sulfate, and concentrated. Column chromatography (25% ethyl acetate/hexanes) gave 20.4 mg (13%) as a light yellow oil. Mass spec.: 365.40 (MH)<sup>+</sup>.

10


1-(Methoxycarbonyl)-2-((2-((2-(trimethylsilyl)ethoxy)methyl)-7-methyl-2H-indazol-5-yl)ethyl 4-nitrophenyl carbonate



15 To a solution of methyl 3-((2-((2-(trimethylsilyl)ethoxy)methyl)-7-methyl-2H-indazol-5-yl)-2-hydroxypropanoate (20 mg, 55  $\mu$ moles) in pyridine (1 mL) was added 4-nitrophenylchloroformate (55 mg, 5 equiv). The reaction was stirred at room temperature overnight. The reaction was treated with an additional portion of 4-nitrophenylchloroformate (30 mg, 2.7 equiv) and stirred at room  
20 temperature for 8 hours. The reaction was poured into diethyl ether, washed with 1M potassium bisulfate until very acidic, then saturated bicarbonate, then 1M sodium hydroxide until most of the nitrophenol had been removed, then brine, dried over sodium sulfate, and concentrated to give 50 mg (quant.) of a pale yellow solid which was used immediately in the next reaction. Mass spec.:  
25 530.30 (MH)<sup>+</sup>.

- 70 -

1-(Methoxycarbonyl)-2-(2-((2-(trimethylsilyl)ethoxy)methyl)-7-methyl-2H-indazol-5-yl)ethyl 4-(1,2-dihydro-2-oxoquinazolin-3(4H)-yl)piperidine-1-carboxylate




5

A flask was charged with 1-(methoxycarbonyl)-2-(2-((2-(trimethylsilyl)ethoxy)methyl)-7-methyl-2H-indazol-5-yl)ethyl 4-nitrophenyl carbonate (27 mg, 51  $\mu$ moles) and 3,4-dihydro-3-(piperidin-4-yl)quinazolin-2(1H)-one (34 mg, 2.8 equiv). The solids were dissolved in dimethylformamide (1 mL) and treated with diisopropylethylamine (0.1 mL, 11 equiv). The reaction was stirred at room temperature for 2 d. The reaction was concentrated, dissolved in ethyl acetate, washed with 20% potassium hydroxide (3X), then brine, dried over magnesium sulfate, and concentrated. Column chromatography (100% ethyl acetate) removed baseline material to give 50 mg (quant.). Mass spec.: 622.50 (MH)<sup>+</sup>.

15

1-(Methoxycarbonyl)-2-(7-methyl-1H-indazol-5-yl)ethyl 4-(1,2-dihydro-2-oxoquinazolin-3(4H)-yl)piperidine-1-carboxylate

- 71 -




1-(Methoxycarbonyl)-2-((2-((2-(trimethylsilyl)ethoxy)methyl)-7-methyl-2H-indazol-5-yl)ethyl 4-(1,2-dihydro-2-oxoquinazolin-3(4H)-yl)piperidine-1-carboxylate (50 mg, 40  $\mu$ moles) was dissolved in trifluoroacetic acid (50% in dichloromethane, 5 mL). After 2 h at room temperature, the reaction was concentrated. Column chromatography (50% ethyl acetate/hexanes  $\rightarrow$  100% ethyl acetate) gave 14.5 mg (37%). Mass spec.: 492.15 (MH)<sup>+</sup>.

10

**Example 1**

**3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(1,2-dihydro-2-oxoquinazolin-3(4H)-yl)piperidine-1-carboxylate**



15

To a solution of 1-(methoxycarbonyl)-2-(7-methyl-1H-indazol-5-yl)ethyl 4-(1,2-dihydro-2-oxoquinazolin-3(4H)-yl)piperidine-1-carboxylate (14.5 mg, 30




- 72 -

μmoles) in methanol (1 mL) was added a solution of lithium hydroxide monohydrate (6.2 mg, 5 equiv) in water (1 mL). The reaction was stirred overnight at room temperature. The reaction was concentrated, dissolved in water, treated with 0.1 mL of 1M hydrochloric acid. A precipitate formed and the  
5 reaction was concentrated to give the crude carboxylic acid which was carried on to the next step without purification. Mass spec.: 478.17 (MH)<sup>+</sup>.

To a solution of the crude acid, 4-piperidinopiperidine (9.9 mg, 2 equiv), and diisopropylethylamine (10 μL, 2 equiv) in dimethylformamide (1 mL) and dichloromethane (1 mL) at 0°C was added PyBOP® (16 mg, 1.05 equiv). The ice  
10 bath was removed and stirring continued for 1 h. The reaction was concentrated and purified by preparative HPLC to give the title compound (16 mg, 73%) as its trifluoroacetic acid salt. <sup>1</sup>H-NMR (CD<sub>3</sub>OD) δ 0.10 (m, 0.5), 1.11 (m, 0.5H), 1.50-2.35 (m, 13H), 2.45-3.30 (m, 10H), 3.38 (m, 2H), 3.55 (m, 2H), 4.15-4.70 (m, 6H), 4.85 (m, 1H), 5.76 (m, 1H), 6.98 (d, *J*=7.6, 1H), 7.14 (dd, *J*=7.6, 7.3,  
15 1H), 7.20-7.55 (m, 3H), 7.71 (m, 1H), 8.25 (m, 1H). Mass spec.: 628.29 (MH)<sup>+</sup>.

7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2-H-indazole-5-carbaldehyde




20 To a solution of 7-methyl-1H-indazole-5-carbaldehyde (5.0 g, 31.25 mmol) and N-methyl-dicyclohexylamine (13.5 mL, 62.35 mmol) in dry tetrahydrofuran (120 mL) at 0°C, was added 2-(trimethylsilyl)ethoxymethyl chloride (6.65 mL, 39.5 mmol). The icebath was removed and stirring continued for 5 h. The reaction mixture was diluted with ethyl acetate, washed with water  
25 (2X), brine (2X), dried over sodium sulfate, and concentrated *in vacuo*. Column

- 73 -

chromatography afforded 8.5 g (93%).  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , 300 MHz)  $\delta$  -0.04 (s, 9H), 0.83-1.01 (m, 2H), 2.60 (s, 3H), 3.22-3.34 (m, 2H), 3.60-3.76 (m, 2H), 5.58 (s, 2H), 7.54 (s, 1H), 8.23 (s, 1H), 8.64 (s, 1H), 9.91 (s, 1H). Mass spec.: 291.33 (MH) $^+$ .


5

2-Acetoxy-2-(diethoxyphosphoryl)acetic acid



Glyoxylic acid monohydrate (4.0 g, 43.45 mmol) was suspended in  
10 diethyl phosphite (5.59 mL, 1.0 equiv), warmed to 60°C, and held there for 5 h. The reaction was cooled, diluted with dichloromethane (40 mL), and treated with pyridine (3.51 mL, 1.0 equiv) and acetyl chloride (3.09 mL, 1.0 equiv). A significant exotherm was noted. The reaction was stirred at room temperature for 2 h. The reaction was washed with 1 M hydrochloric acid (2 X 20 mL), then  
15 saturated sodium bicarbonate. The organics were dried over magnesium sulfate, and concentrated to give <2 g as an oil. The aqueous washes were combined and extracted with dichloromethane (4X). The organics were dried over magnesium sulfate and concentrated to give 5.85 g (53%) as an oil which solidified upon standing.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  1.36 (t,  $J=7.0$ , 6H), 2.21 (s, 3H), 4.28  
20 (m, 4H), 5.54 (d,  $J=17.7$ , 1H), 8.90 (bs, 1H). Mass spec.: 255.10 (MH) $^+$ .


Methyl 2-acetoxy-2-(diethylphosphoryl)acetate



- 74 -

To a heterogeneous mixture of 5 M sodium hydroxide (50 mL) and diethyl ether (100 mL) in a fire-polished erlenmeyer flask at 0°C was added *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine (6.37 g, 43.3 mmol) in small portions with swirling (no stirbar). After addition was complete, the mixture was allowed to stand at 0°C for 15 min with occasional swirling. The ethereal was transferred in portions to a suspension of 2-acetoxy-2-(diethoxyphosphoryl)acetic acid (5.50 g, 21.6 mmol) in ether (ca. 50 mL) until the solid had all dissolved and a yellow color persisted. The reaction was allowed to rest at 0°C for 15 min before bubbling nitrogen through the solution to remove unreacted diazomethane. The reaction was concentrated to give 5.90 g (quant.) as a faint yellow oil. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 1.36 (td, *J*=7.0, 2.4, 6H), 2.21 (s, 3H), 3.82 (s, 3H), 4.23 (m, 4H), 5.43 (d, *J*=16.8, 1H). Mass spec.: 269.17 (MH)<sup>+</sup>.

Methyl 2-acetoxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)acrylate




To a solution of methyl 2-acetoxy-2-(diethylphosphoryl)acetate (923 mg, 3.44 mmol) in tetrahydrofuran (7 mL) was added lithium chloride (146 mg, 3.44 mmol). The reaction was stirred until dissolution was complete. The reaction was cooled to -78°C, and treated with *N,N,N',N'*-tetramethylguanidine (0.43 mL, 3.44 mmol) to give a white suspension which was stirred for 10 min. To this was added the 7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2-H-indazole-5-carbaldehyde (1.00 g, 3.44 mmol) in one portion. The reaction was stirred for 1 h at -78°C, and then allowed to slowly warm to room temperature in the dewar. Allowed to stir overnight at room temperature. The reaction was poured into

- 75 -

water/ether. The mixture was extracted with diethyl ether (2X), which were washed with water, then brine, dried over magnesium sulfate, and concentrated. Column chromatography (12% to 25% ethyl acetate/hexanes) gave 825 mg (59%) as a mixture of *Z*- and *E*- isomers as a colorless oil. Major (*Z* isomer): <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ -0.02 (s, 9H), 0.95 (t, J=8.5, 2H), 2.25 (s, 3H), 2.62 (s, 3H), 3.66 (m, 2H), 3.71 (s, 3H), 5.75 (s, 2H), 6.88 (s, 1H), 7.09 (s, 1H), 7.73 (s, 1H), 8.11 (s, 1H). Mass spec.: 405.17 (MH)<sup>+</sup>. Minor (*E* isomer): <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ -0.02 (s, 9H), 0.95 (t, J=8.5, 2H), 2.36 (s, 3H), 2.62 (s, 3H), 3.66 (m, 2H), 3.85 (s, 3H), 5.74 (s, 2H), 7.32 (s, 1H), 7.38 (s, 1H), 7.78 (s, 1H), 8.14 (s, 1H). Mass spec.: 405.17 (MH)<sup>+</sup>.


(*R*)-Methyl 2-acetoxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)propanoate



To a solution of methyl 2-acetoxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)acrylate (825 mg, 2.04 mmol) in degassed (by bubbling nitrogen) dichloromethane (20.00 mL) under nitrogen was added solid (-)-1,2-bis-((2*R*,5*R*)-2,5-diethylphospholano)benzene-(cyclooctadiene)rhodium(I) tetrafluoroborate (100.00 mg), all at once. The reaction was placed under a hydrogen atmosphere (55 psi), and shaken for 6 h. The reaction was concentrated and purified by column chromatography (25% ethyl acetate/hexanes) to give 700 mg (84%) as a colorless oil. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ -0.03 (s, 9H), 0.94 (t, J=8.2, 2H), 2.07 (s, 3H), 2.61 (s, 3H), 3.11 (dd, J=14.3, 8.9, 1H), 3.20 (dd, J=14.3, 4.6, 1H), 3.64 (t, J=8.5, 2H), 3.72 (s, 3H), 5.26 (dd, J=8.5, 4.6, 1H), 5.72 (s, 2H), 6.93 (s, 1H), 7.33 (s, 1H), 8.02 (s, 1H).

- 76 -

(R)-2-Hydroxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)propanoic acid




5

To a solution of (R)-methyl 2-acetoxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)propanoate (700 mg, 1.72 mmol) in tetrahydrofuran (6 mL) and methanol (6 mL) at 0°C was added a solution of lithium hydroxide monohydrate (289 mg, 6.89 mmol) in water (6 mL). The reaction was stirred at 0°C for 1h. The reaction was concentrated, dissolved in 5 mL of water, cooled to 0°C, and treated with 1 M hydrochloric acid until mildly acidic. A non-solid ppt formed. The suspension was extracted with ethyl acetate (2X), which were washed with brine, dried over magnesium sulfate, and concentrated to give 620 mg (quant.) which was pure by LC/MS and was used without purification. Mass spec.: 351.13 (MH)<sup>+</sup>.

15

2-(Methoxymethyl)-7-methyl-2H-indazole-5-carbaldehyde



20

To a solution of 7-methylindazole-5-carboxaldehyde (8.80 g, 54.9 mmol) and N-methyl-dicyclohexylamine (23.6 mL, 110 mmol) in tetrahydrofuran (200 mL) at 0°C was added chloromethyl methyl ether (7.50 mL, 1.8 equiv). The reaction was allowed to gradually warm to room temperature overnight. The reaction was concentrated, dissolved in diethyl ether, washed with water, then 1


- 77 -

M hydrochloric acid, then water, then brine, dried over magnesium sulfate, and concentrated to give an oil. The oil was dissolved in ethyl acetate and treated with hexanes until lasting turbidity. The suspension was heated until a clear solution was obtained and the flask placed in the freezer. The resulting

5 crystalline solid was crushed with a spatula to break it up, reheated to dissolve some of the solids, and placed in the freezer. The solids were filtered, washed with very cold diethyl ether (-78°C), and air-dried to give 5.43 g. The mother liquor was concentrated, redissolved in diethyl ether (ca. 20 mL), cooled to -78°C, and treated with a seed crystal of the product. After 1 h, the resulting

10 solids were filtered, washed with cold diethyl ether (-78°C), and air-dried to give an additional 1.05 g (total yield = 58%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 2.66 (s, 3H), 3.44 (s, 3H), 5.73 (s, 2H), 7.59 (s, 1H), 8.09 (s, 1H), 8.32 (s, 1H), 9.97 (s, 1H). Mass spec.: 205.19 (MH)<sup>+</sup>.

15 Methyl 2-acetoxy-3-(2-(methoxymethyl)-7-methyl-2H-indazol-5-yl)acrylate



To a solution of methyl 2-acetoxy-2-(diethylphosphoryl)acetate (4.89 g, 18.2 mmol) in tetrahydrofuran (25 mL) was added lithium chloride (0.74 g, 17.5


20 mmol). The reaction was stirred until dissolution was complete. The reaction was cooled to -78°C, and treated with tetramethylguanidine (2.20 mL, 17.5 mmol) to give a white suspension which was stirred for 10 min. To this was added 2-(methoxymethyl)-7-methyl-2H-indazole-5-carbaldehyde (3.10 g, 15.2 mmol) in one portion. After 10 min, the ice bath was concentrated and the

25 reaction stirred overnight. The reaction was poured onto water / diethyl ether, and the layers separated. The ethereal was washed with water, then brine, dried over magnesium sulfate, and concentrated. Column chromatography gave

- 78 -

recovered 2-(methoxymethyl)-7-methyl-2H-indazole-5-carbaldehyde (0.57 g, 18%) and the title compound (2.86 g, 59%) as a mixture of *Z*- and *E*- isomers as a colorless oil. Major (*Z* isomer): <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 2.25 (s, 3H), 2.62 (s, 3H), 3.40 (s, 3H), 3.71 (s, 3H), 5.69 (s, 2H), 6.88 (s, 1H), 7.09 (s, 1H), 7.72 (s, 1H), 8.10 (s, 1H). Mass spec.: 319.18 (MH)<sup>+</sup>. Minor (*E* isomer): <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 2.35 (s, 3H), 2.62 (s, 3H), 3.40 (s, 3H), 3.85 (s, 3H), 5.69 (s, 2H), 7.32 (s, 1H), 7.38 (s, 1H), 7.78 (s, 1H), 8.14 (s, 1H). Mass spec.: 319.18 (MH)<sup>+</sup>.


10 (R)-Methyl 2-acetoxy-3-(2-(methoxymethyl)-7-methyl-2H-indazol-5-yl)propanoate



A solution of methyl 2-acetoxy-3-(2-(methoxymethyl)-7-methyl-2H-indazol-5-yl)acrylate (2.80 g, 8.8 mmol) in dichloromethane (20 mL) was degassed by passing a stream of nitrogen through the solution. To this solution was quickly added (-)-1,2-bis((2*R*,5*R*)-2,5-diethylphospholano)benzene(cyclooctadiene) rhodium (I) trifluoromethylsulfonate (100 mg, 0.016 equiv) as a solid. The reaction was placed under a hydrogen atmosphere (55 psi) and shaken overnight. The reaction was concentrated and purified by column chromatography (50% ethyl acetate/hexanes) to give 2.74 g (97%) as a colorless oil. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 2.08 (s, 3H), 2.61 (s, 3H), 3.11 (dd, *J*=14.3, 8.9, 1H), 3.20 (dd, *J*=14.3, 4.6, 1H), 3.39 (s, 3H), 3.72 (s, 3H), 5.26 (dd, *J*=8.9, 4.6, 1H), 5.68 (s, 2H), 6.93 (s, 1H), 7.33 (s, 1H), 8.02 (s, 1H).

(R)-2-Hydroxy-3-(2-(methoxymethyl)-7-methyl-2H-indazol-5-yl)propanoic acid


- 79 -



To a solution of (R)-methyl 2-acetoxy-3-(2-(methoxymethyl)-7-methyl-2H-indazol-5-yl)propanoate (2.70 g, 8.4 mmol) in tetrahydrofuran (20 mL) and methanol (20 mL) at 0°C was added a solution of lithium hydroxide monohydrate (1.41 g, 4.0 equiv) in water (20 mL). The reaction was stirred at 0°C for 1 h. The reaction was concentrated, dissolved in water (5 mL), cooled to 0°C, and treated with 1 M hydrochloric acid until mildly acidic. The solution was extracted extensively with ethyl acetate and then dichloromethane. The organics were combined, dried over magnesium sulfate, and concentrated to give 1.40 g (63%) as an oil which solidified to a crystalline solid upon standing. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 2.40 (s, 3H), 2.78 (dd, *J*=14.0, 7.9, 1H), 3.00 (dd, *J*=14.0, 4.0, 1H), 3.18 (s, 3H), 4.24 (dd, *J*=7.9, 4.3, 1H), 5.47 (s, 2H), 6.85 (s, 1H), 7.22 (s, 1H), 7.90 (s, 1H). Mass spec.: 265.08 (MH)<sup>+</sup>.

15

(R)-Methyl 2-hydroxy-3-(2-(methoxymethyl)-7-methyl-2H-indazol-5-yl)propanoate



20


To a heterogeneous mixture of 5 M sodium hydroxide (20 mL) and diethyl ether (60 mL) in a fire-polished erlenmeyer flask at 0°C was added N-methyl-N'-nitro-N-nitrosoguanidine (1.17 g, 7.95 mmol) in small portions with swirling (no stirbar). After addition was complete, the mixture was allowed to



- 80 -

stand at 0°C for 15 min with occasional swirling. The ethereal was transferred in portions to a suspension of (R)-2-hydroxy-3-(2-(methoxymethyl)-7-methyl-2H-indazol-5-yl)propanoic acid (1.40 g, 5.30 mmol) in dichloromethane (20 mL) until the solid had all dissolved and a yellow color persisted. The reaction was  
5 allowed to rest at room temperature for ca. 5 min. before bubbling nitrogen through the solution to remove unreacted diazomethane. The reaction was concentrated and purified by column chromatography (50% to 75% ethyl acetate/hexanes) to give 1.47 g (100%) as a colorless oil. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 1.60 (bs, 1H), 2.58 (s, 3H), 2.95 (dd, *J*=13.9, 7.0, 1H), 3.14 (dd, *J*=13.9,  
10 4.0, 1H), 3.36 (s, 3H), 3.76 (s, 3H), 4.46 (bm, 1H), 5.65 (s, 2H), 6.90 (s, 1H), 7.31 (s, 1H), 7.99 (s, 1H). Mass spec.: 279.11 (MH)<sup>+</sup>.

2-(Benzoyloxy)-2-(diethoxyphosphoryl)acetic acid




15

Glyoxylic acid monohydrate (20.10 g, 218 mmol) was suspended in diethyl phosphite (28.1 mL, 1.0 equiv) and warmed to 60°C, and held there for 5 h. The reaction was cooled to 0°C, diluted with dichloromethane (200 mL), and treated with pyridine (17.7 mL, 1.0 equiv) and benzoyl chloride (25.3 mL, 1.0  
20 equiv). The ice bath was removed and stirring continued overnight. The reaction was concentrated, diluted with ethyl acetate, washed with water, then 1 M potassium bisulfate, then brine, dried over magnesium sulfate, and concentrated to give an oil. The oil was triturated with ether to give a white powder which was filtered, washed with diethyl ether, and air dried to give 29.0 g (42%). <sup>1</sup>H-NMR  
25 (CDCl<sub>3</sub>, 500 MHz) δ 1.35 (m, 6H), 4.31 (m, 4H), 5.82 (d, *J*=17.7, 1H), 7.46 (m, 2H), 7.60 (m, 1H), 7.94 (bs, 1H), 8.12 (m, 2H).


- 81 -

## 1-(Diethoxyphosphoryl)-2-methoxy-2-oxoethyl benzoate



Diazomethane was generated in 3 fire-polished erlenmeyer flasks using  
 5 1/3 of each N-methyl-N'-nitro-N-nitrosoguanidine (17.1 g 116. mmol), 5 M  
 sodium hydroxide (200 mL), and diethyl ether (450 mL) at 0°C by adding the  
 guanidine in small portions with swirling to the other two. After addition was  
 complete, the mixture was allowed to stand at 0°C for 10 min with occasional  
 swirling. The ethereal was transferred in portions to a suspension of 2-  
 10 (benzoyloxy)-2-(diethoxyphosphoryl)acetic acid (21.0 g, 66.4  
 mmol) in dichloromethane (ca. 20 mL) until the solid had all dissolved and a  
 yellow color persisted. The reaction was allowed to rest at 0°C for 15 min, before  
 bubbling nitrogen through the solution to remove most of the unreacted  
 diazomethane (reaction went almost colorless). The reaction was concentrated to  
 15 give the 22.0 g (quant.) as a faint yellow oil which was used without purification.  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 1.37 (m, 6H), 3.85 (s, 3H), 4.28 (m, 4H), 5.71 (d,  
 J=16.8, 1H), 7.47 (m, 2H), 7.61 (m, 1H), 8.11 (m, 2H).

1-Methoxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-  
 20 oxoprop-2-en-2-yl benzoate




- 82 -

To a solution of 1-(diethoxyphosphoryl)-2-methoxy-2-oxoethyl benzoate (13.7 g, 41.3 mmol) in tetrahydrofuran (70 mL) was added lithium chloride (1.75 g, 41.3 mmol). The reaction was stirred until dissolution was complete. The reaction was cooled to -78°C, and treated with N,N,N',N'-tetramethylguanidine (5.20 mL, 41.3 mmol) to give a white suspension which was stirred for 10 min. To this was added 7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazole-5-carbaldehyde (10.0 g, 34.4 mmol) in one portion. The reaction was stirred for 10 min at -78°C, placed in a 0°C bath, and allowed to slowly warm to room temperature overnight. The reaction was diluted with ethyl acetate, washed with water, then 1 M potassium bisulfate, then saturated sodium bicarbonate, then brine, dried over magnesium sulfate, and concentrated. Column chromatography (25% ethyl acetate/hexanes) gave 15.10 g (94%) as a mixture of *Z*- and *E*-isomers as a viscous colorless oil. Major (*Z* isomer): <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 0.02 (s, 9H), 0.95 (m, 2H), 2.63 (s, 3H), 3.66 (m, 2H), 3.73 (s, 3H), 5.74 (s, 2H), 7.02 (s, 1H), 7.15 (s, 1H), 7.51 (m, 2H), 7.64 (m, 1H), 7.80 (s, 1H), 8.12 (s, 1H), 8.17 (m, 2H). Mass spec.: 467.18 (MH)<sup>+</sup>. Minor (*E* isomer): <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 0.04 (s, 9H), 0.92 (m, 2H), 2.50 (s, 3H), 3.62 (m, 2H), 3.85 (s, 3H), 5.69 (s, 2H), 7.37 (s, 1H), 7.49 (s, 1H), 7.54 (m, 2H), 7.67 (m, 1H), 7.82 (s, 1H), 8.07 (s, 1H), 8.25 (m, 2H). Mass spec.: 467.18 (MH)<sup>+</sup>.

20


(*R*)-1-Methoxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxopropan-2-yl benzoate



- 83 -

A Parr shaker was purged with nitrogen. A flask containing 1-methoxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxoprop-2-en-2-yl benzoate (15.7 g, 33.7 mmol) was purged with nitrogen. The substrate was dissolved in dichloromethane (100 mL). The solution was transferred via canula  
 5 into the Parr shaker and degassed by passing a stream of nitrogen into the solution for 30 min. (-)-1,2-Bis-((2R,5R)-2,5-diethylphospholano)benzene (cyclooctadiene)rhodium(I)tetrafluoroborate (300 mg) was quickly poured into the flask which was re-sealed and purged for an additional 5 min. The Parr shaker was pressurized to 60 psi of hydrogen and shaken overnight. The reaction  
 10 was concentrated and purified by column chromatography to give 15.5 g (98%) as a colorless oil. Mass spec.: 469.24 (MH)<sup>+</sup>.

(R)-Methyl 2-hydroxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)propanoate



15


To a solution of (R)-1-methoxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxopropan-2-yl benzoate (14.2 g, 30.3 mmol) in tetrahydrofuran (80 mL) and methanol (80 mL) at 0°C was  
 20 added a solution of lithium hydroxide monohydrate (5.09 g, 121 mmol) in water (80 mL). The reaction was stirred at 0°C for 1 h. The ice bath was removed and stirring continued at room temperature for 2 h. The reaction was concentrated, dissolved in water (5 mL), cooled to 0°C, and treated with 1 M hydrochloric acid until mildly acidic. A non-solid ppt formed. The suspension was extracted with  
 25 ethyl acetate (3X), which were washed with brine, dried over magnesium sulfate, and concentrated to give the crude hydroxy acid as an oil with white solid

- 84 -

forming in it (probably residual benzoic acid) which was used without purification. Mass spec.: 349.34 (M-H)<sup>-</sup>.

To a heterogeneous solution of 5 M sodium hydroxide (150 mL) and diethyl ether (450 mL) in a fire-polished erlenmeyer flask at 0°C was added N-methyl-N'-nitro-N-nitrosoguanidine (13.4 g, 90.9 mmol) in small portions with swirling (no stirbar). After addition was complete, the mixture was allowed to stand at 0°C for 15 min with occasional swirling. The ethereal was transferred in portions to a solution of the crude hydroxy acid prepared above in a minimum of diethyl ether until a yellow color persisted. The reaction was allowed to rest at room temperature for 5 min, before bubbling nitrogen through the solution to remove most of the unreacted diazomethane (reaction went almost colorless). The reaction was concentrated and purified by column chromatography (25 to 50% ethyl acetate/hexanes) to give 10.5 g (95%) of the title compound as a colorless oil. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ -0.03 (s, 9H), 0.94 (m, 2H), 2.60 (s, 3H), 2.69 (d, J=6.1, 1H), 2.98 (dd, J=14.0, 7.0, 1H), 3.16 (dd, J=13.7, 4.3, 1H), 3.63 (m, 2H), 3.78 (s, 3H), 4.49 (m, 1H), 5.71 (s, 2H), 6.92 (s, 1H), 7.33 (s, 1H), 8.01 (s, 1H). Mass spec.: 365.02 (MH)<sup>+</sup>.

(R)-2-Hydroxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)propanoic acid (610 mg, 1.74 mmol), diisopropylethylamine (0.61 mL, 3.48 mmol), and 4-piperidinopiperidine (585.76 mg, 2.0 equiv) in dimethylformamide (20 mL) and dichloromethane (20




To a solution of the (R)-2-hydroxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)propanoic acid (610 mg, 1.74 mmol), diisopropylethylamine (0.61 mL, 3.48 mmol), and 4-piperidinopiperidine (585.76 mg, 2.0 equiv) in dimethylformamide (20 mL) and dichloromethane (20

- 85 -

mL) at 0°C was added PyBOP® (951 mg, 1.83 mmol). The reaction was stirred at 0°C for 2 h. The reaction was concentrated, diluted with ethyl acetate, washed with water, then brine, dried over magnesium sulfate, and concentrated. The product was purified by column chromatography (4:96:1  
 5 methanol/dichloromethane/triethylamine) to give 795 mg (91%). Mass spec.: 501.40 (MH)<sup>+</sup>.

(R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-nitrophenyl carbonate




10

To a solution of (R)-2-hydroxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-1-one (795 mg, 1.59 mmol) and diisopropylethylamine (0.55 mL, 3.18  
 15 mmol) in dichloromethane (10 mL) at 0°C was added 4-nitrophenyl-chloroformate (352 mg, 1.10 equiv) followed by dimethylaminopyridine (10 mg). The ice bath was removed and stirring continued for 7 h. The reaction was treated with an additional portion of diisopropylethylamine (0.25 mL, 1.45 mmol), 4-nitrophenyl-chloroformate (352 mg, 1.10 equiv), and  
 20 dimethylaminopyridine (10 mg) and stirred overnight at room temperature. The reaction was concentrated, dissolved in ethyl acetate, washed with saturated sodium bicarbonate (3X), then brine, dried over magnesium sulfate, and concentrated to give 1.06 g (quant.) as a light brown oil which was used without purification. Mass spec.: 666.31 (MH)<sup>+</sup>.

25

- 86 -

(R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinazolin-3(4H)-yl)piperidine-1-carboxylate



5


To a solution of (R)-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-nitrophenyl carbonate (150 mg, 0.23 mmol) and 3-(piperidin-4-yl)-3,4-dihydroquinazolin-2(1H)-one (78.2 mg, 1.50 equiv) in dimethylformamide (1 mL) was added diisopropylethylamine (79  $\mu$ L). The reaction was stirred at room temperature overnight. The reaction was concentrated and purified by column chromatography to give 90 mg (53%) as a colorless film. Mass spec.: 758.56 (MH)<sup>+</sup>.

15

### Example 2

(R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinazolin-3(4H)-yl)piperidine-1-carboxylate

- 87 -




(R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinazolin-3(4H)-yl)piperidine-1-carboxylate (90 mg, 0.12 mmol) was dissolved in trifluoroacetic acid (50% in dichloromethane, 6 mL) and stirred at room temperature for 2 h. The reaction was concentrated and purified by column chromatography (5:95:1 methanol/dichloromethane/triethylamine). The compound was then dissolved in 10% methanol/dichloromethane and passed through a plug of basic alumina to give 23 mg (31%) as a white solid. <sup>1</sup>H-NMR (CD<sub>3</sub>OD, 500 MHz) δ -0.21 (m, 0.7H), 0.74 (m, 1H), 1.10-2.05 (m, 17H), 2.15-2.60 (m, 7H), 2.83 (m, 3H), 3.07 (m, 1.5 H), 3.16 (dd, J=12.8, 5.8, 0.7H), 3.70-4.55 (m, 7H), 5.42 (m, 0.3H), 5.50 (dd, J=9.5, 6.1, 0.7 H), 6.70 (m, 1H), 6.85 (m, 1H), 6.90-7.15 (m, 3H), 7.40 (bs, 1H), 7.93 (bs, 1H). Mass spec.: 628.34 (MH)<sup>+</sup>.

15

(R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(8-fluoro-2-oxo-1,2-dihydroquinazolin-3(4H)-yl)piperidine-1-carboxylate



- 88 -




To a solution of (R)-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-nitrophenyl carbonate (150 mg, 0.23 mmol) and 8-fluoro-3-(piperidin-4-yl)-3,4-dihydroquinazolin-2(1H)-one (84.2 mg, 1.50 equiv) in dimethylformamide (1 mL) was added diisopropylethylamine (79  $\mu$ L, 0.45 mmol). The reaction was stirred at room temperature overnight. The reaction was concentrated and purified by column chromatography to give 78 mg (45%) as a colorless film.

Mass spec.: 776.58 (MH)<sup>+</sup>.

### Example 3


(R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(8-fluoro-2-oxo-1,2-dihydroquinazolin-3(4H)-yl)piperidine-1-carboxylate



- 89 -

(R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(8-fluoro-2-oxo-1,2-dihydroquinazolin-3(4H)-yl)piperidine-1-carboxylate (78 mg, 0.10 mmol) was dissolved in trifluoroacetic acid (50% in dichloromethane, 6 mL) and stirred at room temperature for 2 h. The reaction was concentrated and purified by column chromatography (5:95:1 methanol/dichloromethane/triethylamine). The residue was dissolved in 10% methanol/dichloromethane and passed through a plug of basic alumina to give 40.1 mg (62%) as a white solid. <sup>1</sup>H-NMR (CD<sub>3</sub>OD, 500 MHz) δ -0.20-0.30 (m, 0.6H), 0.80-1.20 (m, 0.7H), 1.40-2.35 (m, 16H), 2.45-2.83 (m, 6H), 2.90-3.48 (m, 5H), 4.00-4.83 (m, 7H), 5.71 (m, 0.3 H), 5.79 (dd, J=9.5, 6.1, 0.7H), 7.00-7.25 (m, 3H), 7.25-7.41 (m, 1H), 7.69 (bs, 1H), 8.22 (bs, 1H). Mass spec.: 646.50 (MH)<sup>+</sup>.

(R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate




To a solution of (R)-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-nitrophenyl carbonate (150 mg, 0.23 mmol) and 3-(piperidin-4-yl)quinolin-2(1H)-one hydrochloride (89.5 mg, 1.50 equiv) in dimethylformamide (1 mL) was added diisopropylethylamine (0.16 mL, 0.90 mmol), and stirred at room temperature overnight. The reaction was concentrated and purified by column

- 90 -

chromatography to give 102 mg (60%) as a colorless film. Mass spec.: 755.53 (MH)<sup>+</sup>.

#### Example 4


- 5 (R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate



- (R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate (102 mg, 0.14 mmol) was dissolved in trifluoroacetic acid (50% in dichloromethane, 6 mL), and stirred at room temperature for 2 h. The reaction was concentrated and purified by column chromatography (5:95:1 methanol/dichloromethane/triethylamine). The residue was dissolved in 10% methanol/dichloromethane and passed through a plug of basic alumina to give 56.4 mg (67%) as a white solid. <sup>1</sup>H-NMR (CD<sub>3</sub>OD, 500 MHz) δ -0.50 -0.01 (m, 0.7H), 0.50-0.85 (m, 0.8H), 1.00-2.05 (m, 16H), 2.05-2.55 (m, 7H), 2.55-3.00 (m, 4H), 3.00-3.15 (m, 2H), 3.70-4.50 (m, 4H), 5.41 (m, 0.3H), 5.48 (dd, J=9.5, 6.1, 0.6H), 7.01 (m, 1H), 7.11 (m, 1H), 7.21 (m, 1H), 7.36 (m, 2H), 7.44-7.65 (m, 2H), 7.90 (m, 1H). Mass spec.: 625.33 (MH)<sup>+</sup>.

(R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-4-phenyl-2,3-dihydroimidazol-1-yl)piperidine-1-carboxylate


- 91 -



To a solution of (R)-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-nitrophenyl carbonate (150 mg, 0.23 mmol) and 4-phenyl-1-(piperidin-4-yl)-1H-imidazol-2(3H)-one (*PCT Int. Appl.* 1998, WO 9811128 A1) (82.2 mg, 1.50 equiv) in dimethylformamide (1 mL) was added diisopropylethylamine (79  $\mu$ L, 0.45 mmol) and stirred at room temperature overnight. The reaction was concentrated and purified by column chromatography to give 74 mg (43%) as a colorless film. Mass spec.: 770.57 (MH)<sup>+</sup>.

#### Example 5


(R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-4-phenyl-2,3-dihydroimidazol-1-yl)piperidine-1-carboxylate



- 92 -

(R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-4-phenyl-2,3-dihydroimidazol-1-yl)piperidine-1-carboxylate (74 mg, 96.1  $\mu$ mol) was dissolved in trifluoroacetic acid (50% in dichloromethane, 6 mL), and stirred at room temperature for 2 h. The reaction was concentrated and purified by column chromatography (5:95:1 methanol/dichloromethane/triethylamine). The residue was dissolved in 10% methanol/dichloromethane and passed through a plug of basic alumina to give 31.7 (52%) as a white solid.  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , 500 MHz)  $\delta$  -0.50 -0.10 (m, 0.7H), 0.50-0.90 (m, 1H), 1.05-1.55 (m, 12H), 1.55-2.60 (m, 16H), 2.73 (m, 0.8H), 2.85-3.19 (m, 2.6H), 3.30-3.60 (m, 1H), 3.65-4.18 (m, 3.4H), 4.26 (m, 1H), 4.40 (m, 1H), 5.25-5.57 (m, 1H), 6.90-7.10 (m, 1H), 7.10-7.60 (m, 7H), 7.84-7.97 (bs, 1H). Mass spec.: 638.46 (M-H).

*tert*-Butyl 4-hydroxy-4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate



15


A flask was charged with 3-bromoquinolin-2(1H)-one (*U.S. Pat. Appl. Publ.* 2002, US 2002099208 A1; Payack, J.F. *et al. J. Org. Chem.* 2005, 70, 1, 175.) (1.0 g, 4.46 mmol) and sodium hydride (118 mg, 4.91 mmol). The solids were dissolved in tetrahydrofuran (30 mL), stirred for 15 min, and cooled to  $-78^\circ\text{C}$ . The solution was treated with *tert*-butyllithium (1.7 M in pentane, 5.25 mL, 8.93 mmol) and stirred for 1 h. To this was added *N-tert*-butoxycarbonyl-4-piperidone (889 mg, 1.0 equiv). The ice bath was removed and stirring continued for 1 h. The reaction was quenched by addition of saturated ammonium chloride. The reaction was diluted with diethyl ether, washed with water (2X), then brine, dried over magnesium sulfate, and concentrated. The resulting oil was triturated with diethyl ether to give a white solid which was filtered, washed with diethyl

- 93 -

ether, and air dried to give 430 mg (28%) as a white powder.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ , 500 MHz)  $\delta$  1.48 (s, 9H), 1.57 (bs, 2H), 1.87 (m, 2H), 2.13 (m, 2H), 3.37 (m, 2H), 4.06 (m, 2H), 7.28 (m, 2H), 7.53 (m, 2H), 7.60 (m, 1H), 7.69 (s, 1H). Mass spec.: 367.35 ( $\text{MNa}^+$ ).

5


3-(4-Hydroxypiperidin-4-yl)quinolin-2(1H)-one



*tert*-Butyl 4-hydroxy-4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate (300 mg, 0.87 mmol) was dissolved in trifluoroacetic acid (50% in dichloromethane, 6 mL). The reaction was stirred 30 min and concentrated. The residue was dissolved in water, extracted with dichloromethane (2X) which were discarded. The aqueous was made basic with solid potassium carbonate. The resulting solid was filtered to give 145 mg (68%) as a white powder. Mass spec.: 245.35 ( $\text{MH}^+$ ).

15

(R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-hydroxy-4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate



20


- 94 -

To a solution of (R)-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-nitrophenyl carbonate (150 mg, 0.23 mmol) and 3-(4-hydroxypiperidin-4-yl)quinolin-2(1H)-one (55 mg, 1.0 equiv) in dimethylformamide (1 mL) was added  
 5 diisopropylethylamine (79  $\mu$ L, 0.45 mmol). The reaction was stirred at room temperature overnight. The reaction was concentrated and purified by column chromatography (3:97:1 methanol/dichloromethane/triethylamine) to give 90 mg (52%) as a foam solid. Mass spec.: 771.48 (MH)<sup>+</sup>.

10

**Example 6**

**(R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-hydroxy-4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate**




15

(R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-hydroxy-4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate (90 mg, 0.12 mmol) was dissolved in trifluoroacetic acid (50% in dichloromethane, 6 mL) and stirred at room  
 20 temperature for 2 h. The reaction was concentrated and purified by column chromatography (5:95:1 to 10:90:1 methanol/dichloromethane/triethylamine). The residue was dissolved in 10% methanol/dichloromethane and passed through a plug of basic alumina to give 41 mg (55%) as a white solid. <sup>1</sup>H-NMR (CD<sub>3</sub>OD, 500 MHz)  $\delta$  -0.40-0.00 (m, 0.6H), 0.50-0.95 (m, 0.7H), 1.10-2.75 (m, 21H), 2.83  
 25 (m, 0.7H), 3.02-3.65 (m, 6H), 3.68-4.35 (m, 3H), 4.37-4.57 (m, 1H), 5.40-5.65

- 95 -

(m, 1H), 7.00-7.15 (m, 1H), 7.15-7.25 (m, 1H), 7.25-7.35 (m, 1H), 7.40-7.52 (m, 2H), 7.57-7.70 (m, 1H), 7.87-8.03 (m, 1H). Mass spec.: 641.61 (MH)<sup>+</sup>.

(R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(7-fluoro-2-oxo-1,2-dihydroquinazolin-3(4H)-yl)piperidine-1-carboxylate




To a solution of (R)-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-nitrophenyl carbonate (150 mg, 0.23 mmol) and 7-fluoro-3-(piperidin-4-yl)-3,4-dihydroquinazolin-2(1H)-one (56.2 mg, 1.0 equiv) in dimethylformamide (1 mL) was added diisopropylethylamine (79  $\mu$ L, 0.45 mmol) and the reaction stirred at room temperature overnight. The reaction was concentrated and purified by column chromatography (3:97:1 methanol/dichloromethane/triethylamine) to give 93 mg (53%) as a tan foam solid. Mass spec.: 776.48 (MH)<sup>+</sup>.

#### Example 7

(R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(7-fluoro-2-oxo-1,2-dihydroquinazolin-3(4H)-yl)piperidine-1-carboxylate




- 96 -



(R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(7-fluoro-2-oxo-1,2-dihydroquinazolin-3(4H)-yl)piperidine-1-carboxylate (93 mg, 0.12 mmol) was dissolved in trifluoroacetic acid (50% in dichloromethane, 6 mL) and stirred at room temperature for 2 h. The reaction was concentrated and purified by column chromatography (5:95:1 to 10:90:1 methanol/dichloromethane/triethylamine). The residue was dissolved in 10% methanol/dichloromethane and passed through a plug of basic alumina to give 41 mg (53%) as a white powder. <sup>1</sup>H-NMR (CD<sub>3</sub>OD, 500 MHz) δ -0.38-0.15 (m, 0.7H), 0.60-0.95 (m, 1H), 1.18-2.18 (m, 17H), 2.29 (m, 0.7H), 2.35-2.50 (m, 3H), 2.58-3.01 (m, 3.4H), 3.01-3.25 (m, 2.4H), 3.65-4.60 (m, 7H), 5.47 (m, 0.4H), 5.55 (dd, J=9.5, 6.1, 0.7 H), 6.51 (m, 0.8H), 6.62 (m, 0.8H), 6.90-7.20 (m, 2.2H), 7.45 (bs, 1.1H), 7.98 (bs, 1H). Mass spec.: 648.64 (MH)<sup>+</sup>.

(R)-Methyl 3-(2-(methoxymethyl)-7-methyl-2H-indazol-5-yl)-2-((4-nitrophenoxy)carbonyloxy)propanoate


- 97 -



To a solution of (R)-methyl 2-hydroxy-3-(2-(methoxymethyl)-7-methyl-2H-indazol-5-yl)propanoate (1.45 g, 5.21 mmol) and diisopropylethylamine (2.73 mL, 3.0 equiv) in dichloromethane (27 mL) at 0°C was added 4-nitrophenyl-chloroformate (1.58 g, 1.5 equiv) and *N,N*-dimethylaminopyridine (10 mg). The ice bath was removed and stirring continued for 7 h. The reaction was treated with an additional portion of diisopropylethylamine (1.5 mL, 1.65 equiv), 4-nitrophenyl-chloroformate (1.6 g, 1.5 equiv), and *N,N*-dimethylaminopyridine (10 mg) and stirred overnight. The reaction was concentrated, dissolved in ethyl acetate, washed with water, then 1 M potassium bisulfate, then saturated sodium bicarbonate (5X), then brine, dried over magnesium sulfate, and concentrated to give 6.0 g (quant.) as a light brown oil, which was used immediately without purification. Mass spec.: 444.10 (MH)<sup>+</sup>.


(R)-1-Methoxy-3-(2-(methoxymethyl)-7-methyl-2H-indazol-5-yl)-1-oxopropan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate

- 98 -



A flask was charged with (R)-methyl 3-(2-(methoxymethyl)-7-methyl-2H-indazol-5-yl)-2-((4-nitrophenoxy)carbonyloxy)propanoate (2.31 g, 5.20  
 5 mmol), 3-(piperidin-4-yl)quinolin-2(1H)-one (1.78 g, 1.5 equiv), diisopropylethylamine (1.82 mL, 2.0 equiv), and dimethylformamide (20 mL). The reaction was stirred at room temperature for 8 h and concentrated under vacuum. The resulting residue was dissolved in ethyl acetate and washed with  
 10 water to give a suspension which was exhaustively extracted with ethyl acetate then dichloromethane. The organics were dried over magnesium sulfate and concentrated. Column chromatography (25% ethyl acetate/hexanes to 10% methanol/ethyl acetate) gave 2.40 g (86%) as a light yellow foam solid. Mass spec.: 533.30 (MH)<sup>+</sup>.


15 (R)-1-Methoxy-3-(7-methyl-1H-indazol-5-yl)-1-oxopropan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate



- 99 -

To a solution of (R)-1-methoxy-3-(2-(methoxymethyl)-7-methyl-2H-indazol-5-yl)-1-oxopropan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate (1.20 g, 2.25 mmol) in methanol (20 mL) was added acetyl chloride (0.40 mL, 5.62 mmol). The reaction was warmed to reflux and held there for 1 h. The reaction was concentrated by rotary evaporation under high vacuum and purified by column chromatography (5% methanol/dichloromethane) to give 1.09 g (99%) as a white foam solid. Mass spec.: 489.29 (MH)<sup>+</sup>.

(R)-3-(7-Methyl-1H-indazol-5-yl)-2-(4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxyloxy)propanoic acid




To a solution of (R)-1-methoxy-3-(7-methyl-1H-indazol-5-yl)-1-oxopropan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate (1.20 g, 2.46 mmol) in methanol (10 mL) and tetrahydrofuran (10 mL) at 0°C was added a precooled solution of lithium hydroxide monohydrate (309 mg, 3.0 equiv) in water (10 mL). After 2 h, the reaction was concentrated under high vacuum (<25°C). The resulting residue was dissolved in water (20 mL), cooled to 0°C, and acidified to ca. pH 2 with 1 N hydrochloric acid. The suspension was maintained at 0°C for 1 h, filtered, and the solid washed with cold water. The resulting white solid was air dried and then dried under high vacuum overnight to give 1.04 g (89%) as a white solid. Mass spec.: 475.30 (MH)<sup>+</sup>.

- 100 -

**Example 4**

**(R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate**




5

To a solution of (R)-3-(7-methyl-1H-indazol-5-yl)-2-(4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxyloxy)propanoic acid (500 mg, 1.05 mmol), diisopropylethylamine (0.37 mL, 2.11 mmol), and 4-piperidinopiperidine (355 mg, 2.0 equiv) in dimethylformamide (4 mL) and dichloromethane (4 mL) at 0°C was added PyBOP® (576 mg, 1.11 mmol). The reaction was stirred at 0°C for 4 h, was concentrated, and purified by column chromatography (95:5:1 to 93:7:1 dichloromethane/methanol/triethylamine). The resulting residue was dissolved in 5% methanol/dichloromethane and passed through a basic alumina column to give 600 mg (88%) as a white powder. Mass spec.: 625.56 (MH)<sup>+</sup>.

15

*tert*-Butyl 4-(5,6-dihydropyridin-1(2H)-yl)piperidine-1-carboxylate




4-Oxo-piperidine-1-carboxylic acid *tert*-butyl ester (797 mg, 4 mmol), 1,2,3,6-tetrahydropyridine (349 mg, 4.2 mmol), sodium cyanoborohydride (126 mg, 2 mmol), zinc chloride (410 mg, 3.2 mmol), and anhydrous methanol (20

20

- 101 -

mL) were mixed together and the mixture stirred overnight at room temperature. The solvent was evaporated from the mixture and the residue partitioned between 1 N sodium hydroxide and dichloromethane. The phases were separated and the aqueous layer extracted with dichloromethane. The organic extracts were  
5 combined, dried over magnesium sulfate, and the solvent evaporated. The residue was purified by flash column chromatography (1:1 hexanes/ethyl acetate to ethyl acetate to 1:3 methanol/ethyl acetate) to give 800 mg (75%) as a colorless oil.  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>, 400 MHz) δ 5.75–5.70 (m, 1H), 5.68–5.64 (m, 1H), 4.13–4.07 (m, 2H), 3.08 (m, 2H), 2.71–2.60 (m, 4H), 2.48–2.41 (m, 1H), 2.15 (m, 2H),  
10 1.82–1.79 (m, 2H), 1.49–1.38 (m, 11H).

1-(Piperidin-4-yl)-1,2,3,6-tetrahydropyridine




15 *tert*-Butyl 4-(5,6-dihydropyridin-1(2H)-yl)piperidine-1-carboxylate (800 mg, 3 mmol) was dissolved in dichloromethane (54 ml). To this was added trifluoroacetic acid (7.9 ml) and triethyl silane (1.2 ml). The mixture was stirred at room temperature for 3 h. Solvent was removed from the mixture *en vacuo*. The residue was dissolved in saturated sodium bicarbonate (50 ml) and stirred for  
20 30 min. Sodium hydroxide (50 ml, 50% in water) was added to the solution which was then extracted with dichloromethane (3 x 100 ml). The organic extract was dried over sodium sulfate and the solvent evaporated to give 370 mg (74%) as a colorless oil. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 400 MHz) δ 5.74–5.70 (m, 1H), 5.68–5.64 (m, 1H), 3.14–3.07 (m, 4H), 2.64–2.53 (m, 4H), 2.43–2.37 (m, 1H), 2.15 (m, 2H),  
25 1.85–1.82 (m, 2H), 1.73 (s, 1H), 1.47–1.38 (m, 2H).

- 102 -

## Example 8

(R)-1-(4-(5,6-Dihydropyridin-1(2H)-yl)piperidin-1-yl)-3-(7-methyl-1H-indazol-5-yl)-1-oxopropan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate




5

To a solution of (R)-3-(7-methyl-1H-indazol-5-yl)-2-(4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxyloxy)propanoic acid (50 mg, 0.11 mmol), diisopropylethylamine (37  $\mu$ L, 0.21 mmol), and 1-(piperidin-4-yl)-1,2,3,6-tetrahydropyridine (75.7 mg, 2.0 equiv) in dimethylformamide (0.50 mL) and dichloromethane (0.50 mL) at 0°C was added PyBOP® (57.6 mg, 0.11 mmol). The reaction was stirred at 0°C for 4 h. The reaction was concentrated and purified by column chromatography (95:5:1 to 90:10:1 dichloromethane/methanol/triethylamine). The resulting residue was dissolved in 5% methanol/dichloromethane and passed through a basic alumina column to give 35.9 mg (53%) as a white powder. <sup>1</sup>H-NMR (CD<sub>3</sub>OD, 500 MHz)  $\delta$  -0.35-0.15 (m, 0.7H), 0.60-0.93 (m, 1H), 1.15-2.20 (m, 11H), 2.30-3.25 (m, 11H), 3.75-4.58 (m, 4H), 5.35-5.75 (m, 3H), 7.01-7.14 (m, 1H), 7.18 (m, 1H), 7.28 (m, 1H), 7.36-7.50 (m, 2H), 7.53-7.73 (m, 2H), 7.90-8.01 (m, 1H). Mass spec.: 623.33 (MH)<sup>+</sup>.

20

3-Methyl-3,9-diaza-spiro[5.5]undecane dihydrochloride

- 103 -




3-Benzyl-9-methyl-3,9-diaza-spiro[5.5]undecane (Rice, L.M. *et al.* *J. Heterocyclic Chem.* **1964**, 1, 3, 125.) (1.2 g, 4.65 mmol) was dissolved in ethanol (20 mL). To this solution was added hydrochloric acid (4 N in dioxane, 3 mL) and palladium (10% on charcoal, 500 mg). The reaction was shaken on a Parr shaker overnight under 60 psi of hydrogen. The reaction mixture was filtered through a pad of celite, concentrated, and the residue treated with a mixture of ethyl acetate and hexanes, to yield 670 mg (60%) as a white powder.

<sup>1</sup>H-NMR (CD<sub>3</sub>OD) δ 1.68-1.72 (m, 4H), 2.88 (s, 3H), 3.16-3.22 (m, 6H), 3.38 (m, 2H). Mass spec.: 169.12 (MH)<sup>+</sup>.

### Example 9

(R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(9-methyl-3,9-diaza-spiro[5.5]undecan-3-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate



To a solution of (R)-3-(7-methyl-1H-indazol-5-yl)-2-(4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxyloxy)propanoic acid (50 mg, 0.11 mmol), diisopropylethylamine (37 μL, 0.21 mmol), and 3-methyl-3,9-diaza-




- 104 -

spiro[5.5]undecane dihydrochloride (65.2 mg, 2.0 equiv) in dimethylformamide (0.50 mL) and dichloromethane (0.50 mL) at 0°C was added PyBOP® (57.6 mg, 0.11 mmol). The reaction was stirred at 0°C for 4 h. The reaction was concentrated and purified by column chromatography (95:5:1 to 90:10:1  
5 dichloromethane/methanol/triethylamine). The resulting residue was dissolved in 5% methanol/dichloromethane and passed through a basic alumina column to give 52 mg (77%) as a white powder. <sup>1</sup>H-NMR (CD<sub>3</sub>OD, 500 MHz) δ 0.12-0.55 (m, 1H), 0.85 -1.70 (m, 11H), 1.85 (m, 2H), 2.05-2.42 (m, 8H), 2.50 (s, 3H), 2.73-3.30 (m, 8H), 3.65 (bs, 1H), 4.00-4.43 (m, 2H), 5.46 (dd, J=7.6, 7.6, 1H),  
10 7.07 (s, 1H), 7.17 (m, 1H), 7.27 (m, 1H), 7.38-7.46 (m, 2H), 7.50 (bs, 0.5 H), 7.58 (m, 1H), 7.65 (bs, 0.5 H), 7.96 (s, 1H). Mass spec.: 625.33 (MH)<sup>+</sup>.

### Example 10

(R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(piperidin-1-yl)propan-2-yl 4-(2-  
15 oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate




To a solution of (R)-3-(7-methyl-1H-indazol-5-yl)-2-(4-(2-oxo-1,2-  
dihydroquinolin-3-yl)piperidine-1-carboxyloxy)propanoic acid (50 mg, 0.11  
20 mmol), diisopropylethylamine (37 µL, 0.21 mmol), and piperidine (21 µL, 2.0 equiv) in dimethylformamide (0.5 mL) and dichloromethane (0.5 mL) at 0°C was added PyBOP® (57.6 mg, 0.11 mmol). The reaction was stirred at 0°C for 4 h. The reaction was concentrated and purified by column chromatography (95:5:1 to 90:10:1 dichloromethane/methanol/triethylamine). The resulting residue was  
25 dissolved in 5% methanol/dichloromethane and passed through a basic alumina

- 105 -

column to give 55 mg (93%) as a white powder.  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , 500 MHz)  $\delta$  0.75-1.70 (m, 9H), 1.70-1.95 (m, 4H), 2.51 (s, 3H), 2.72-3.04 (m, 3H), 3.11 (m, 4H), 3.19-3.58 (m, 4H), 3.98-4.49 (m, 2H), 5.47 (dd,  $J=7.3$ , 7.3, 1H), 7.07 (s, 1H), 7.17 (m, 1H), 7.27 (m, 1H), 7.43 (m, 2.5H), 7.58 (d,  $J=7.3$ , 1H), 7.65 (bs, 0.5H), 7.96 (s, 1H). Mass spec.: 542.29 (MH) $^+$ .

### Example 11

(R)-1-(4-Cyclohexylpiperazin-1-yl)-3-(7-methyl-1H-indazol-5-yl)-1-oxopropan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate



10


To a solution of (R)-3-(7-methyl-1H-indazol-5-yl)-2-(4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxyloxy)propanoic acid (50 mg, 0.11 mmol), diisopropylethylamine (37  $\mu\text{L}$ , 0.21 mmol), and 1-cyclohexylpiperazine (35.5 mg, 2.0 equiv) in dimethylformamide (0.50 mL) and dichloromethane (0.50 mL) at 0°C was added PyBOP $^{\text{®}}$  (57.6 mg, 0.11 mmol). The reaction was stirred at 0°C for 4 h. The reaction was concentrated and purified by column chromatography (95:5:1 to 90:10:1 dichloromethane/methanol/ triethylamine). The resulting residue was dissolved in 5% methanol/ dichloromethane and passed through a basic alumina column to give 62.3 mg (92%) as a white powder.  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , 500 MHz)  $\delta$  0.75-2.10 (m, 20H), 2.29 (m, 1H), 2.47 (m, 1H), 2.56 (s, 3H), 2.77-3.41 (m, 9H), 3.73 (m, 1H), 4.04-4.49 (m, 2H), 5.52 (dd,  $J=8.2$ , 7.3, 1H), 7.11 (s, 1H), 7.22 (dd,  $J=7.0$ , 7.0, 1H), 7.33 (d,  $J=7.9$ , 1H), 7.42-7.53 (m, 2H), 7.53-7.76 (m, 2H), 8.01 (s, 1H). Mass spec.: 625.33 (MH) $^+$ .

25

- 106 -

## Example 12

(R)-1-(4-(4-Fluorophenyl)piperazin-1-yl)-3-(7-methyl-1H-indazol-5-yl)-1-oxopropan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate



5

To a solution of (R)-3-(7-methyl-1H-indazol-5-yl)-2-(4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxyloxy)propanoic acid (50 mg, 0.11 mmol), diisopropylethylamine (37  $\mu$ L, 0.21 mmol), and N-(4-fluorophenyl)piperazine (38.0 mg, 2.0 equiv) in dimethylformamide (0.50 mL) and dichloromethane (0.50 mL) at 0°C was added PyBOP® (57.6 mg, 0.11 mmol). The reaction was stirred at 0°C for 4 h. The reaction was concentrated and purified by column chromatography (95:5:1 to 90:10:1 dichloromethane/methanol/triethylamine). The resulting residue was dissolved in 5% methanol/dichloromethane and passed through a basic alumina column to give

10 71.2 mg (quant.) as a white powder. <sup>1</sup>H-NMR (CD<sub>3</sub>OD, 500 MHz)  $\delta$  1.25-1.75 (m, 2H), 1.75-2.25 (m, 9H), 2.51 (s, 3H), 2.75 (m, 1H), 2.75-3.30 (m, 13H), 3.48 (m, 3H), 3.80 (m, 1H), 4.05-4.50 (m, 2H), 5.55 (dd, J=7.6, 7.3, 1H), 6.68 (m, 2H), 6.91 (m, 1H), 7.11 (s, 1H), 7.21 (dd, J=7.9, 7.3, 1H), 7.32 (d, J=8.2, 1H), 7.47 (dd, J=7.9, 7.3, 1H), 7.51 (m, 1.5H), 7.61 (d, J=7.6, 1H), 7.67 (bs, 0.5H), 8.00 (s,


15 1H). Mass spec.: 637.28 (MH)<sup>+</sup>.

20

## Example 13


(R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(pyridin-4-yl)piperazin-1-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate

- 107 -



To a solution of (R)-3-(7-methyl-1H-indazol-5-yl)-2-(4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxyloxy)propanoic acid (50 mg, 0.11 mmol), diisopropylethylamine (37  $\mu$ L, 0.21 mmol), and 1-(4-pyridyl)piperazine (34.4 mg, 2.0 equiv) in dimethylformamide (0.50 mL) and dichloromethane (0.50 mL) at 0°C was added PyBOP® (57.6 mg, 0.11 mmol). The reaction was stirred at 0°C for 4 h, was concentrated, and purified by column chromatography (95:5:1 to 90:10:1 dichloromethane/methanol/triethylamine). The resulting residue was dissolved in 5% methanol/dichloromethane and passed through a basic alumina column to give 53.8 mg (80%) as a white powder. <sup>1</sup>H-NMR (CD<sub>3</sub>OD, 500 MHz)  $\delta$  1.25-2.00 (m, 4H), 2.51 (s, 3H), 2.67 (bs, 1H), 2.80-3.31 (m, 7H), 3.35-3.83 (m, 5H), 4.05-4.49 (m, 2H), 5.52 (dd, J=7.6, 7.3, 1H), 6.64 (bs, 2H), 7.12 (s, 1H), 7.22 (dd, J=7.0, 7.0, 1H), 7.32 (d, J=8.2, 1H), 7.47 (m, 1H), 7.52 (m, 1.5H), 7.62 (d, J=7.0, 1H), 7.69 (bs, 0.5H), 7.99 (s, 1H), 8.10 (m, 2H). Mass spec.: 620.28 (MH)<sup>+</sup>.

(±)-2-Hydroxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)propanoic acid



- 108 -


A Parr bottle was charged with 1-methoxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxoprop-2-en-2-yl benzoate (1.0 g, 2.14 mmol), methanol (15 mL), and palladium (10% on charcoal, 100 mg). The Parr shaker was pressurized to 60 psi of hydrogen and shaken for 6 h.

5 The reaction was filtered through celite and concentrated. The resulting residue was dissolved in tetrahydrofuran (8 mL) and methanol (8 mL) and cooled to 0°C. To this was added a solution of lithium hydroxide monohydrate (358 mg, 8.54 mmol) in water (8 mL). The reaction was stirred at 0°C for 1 h, then at room temperature for 2 h. The reaction was concentrated, dissolved in water, cooled to

10 0°C, and acidified with 1 M hydrochloric acid. The resulting mixture was extracted with ethyl acetate (2X). The organics were washed with a minimum of brine, dried over magnesium sulfate, and concentrated. Column chromatography (5% to 10% methanol/dichloromethane) gave 340 mg (45%) as a colorless oil.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ -0.04 (s, 9H), 0.93 (m, 2H), 2.53 (s, 3H), 2.92 (m, 1H), 3.19 (m, 1H), 3.61 (m, 2H), 4.46 (bs, 1H), 5.68 (s, 2H), 6.93 (s, 1H), 7.31 (bs, 1H), 7.94 (bs, 1H). Mass spec.: 351.36 (MH)<sup>+</sup>.

(±)-2-Hydroxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-1-one



20

To a solution of (±)-2-hydroxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)propanoic acid (100 mg, 0.29 mmol), 4-piperidinopiperidine (96.0 mg, 2.0 equiv), and diisopropylethylamine (0.10 mL, 0.57 mmol) in dichloromethane (3.6 mL) at 0°C was added PyBOP<sup>®</sup>


25 (156 mg, 0.30 mmol) in two portions. The reaction was stirred at 0°C for 15

- 109 -

minutes and at room temperature overnight. The reaction was diluted with ethyl acetate, washed with water (2X), then brine, dried over magnesium sulfate, and concentrated. Column chromatography (4% to 10% methanol/dichloromethane) gave 130 mg (91%) as an oil. Mass spec.: 501.36 (MH)<sup>+</sup>.

5

(±)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate



10


To a solution of (±)-2-hydroxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-1-one (130 mg, 0.26 mmol) and diisopropylethylamine (91  $\mu$ L, 0.52 mmol) in dichloromethane (1 mL) at 0°C was added 4-nitrophenyl-chloroformate (57.6 mg, 1.10 equiv). The ice bath was removed and stirring continued for 4 h. The reaction was treated with a solution of 3-(piperidin-4-yl)quinolin-2(1H)-one (88.9 mg, 1.50 equiv) and diisopropylethylamine (91  $\mu$ L, 0.52 mmol) in dimethylformamide (1 mL). The reaction was stirred at room temperature overnight. The reaction was poured into water/dichloromethane. The mixture was extracted with dichloromethane (2X) which was washed with saturated sodium bicarbonate, then water (2X), then brine, dried over magnesium sulfate, and concentrated. Column chromatography (5% to 20% methanol/dichloromethane) gave 112 mg (57%) as an oil.

20

- 110 -

## Example 14

(±)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate




5

(±)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate (112 mg, 0.15 mmol) was dissolved in trifluoroacetic acid (50% in dichloromethane, 6 mL) and stirred at room temperature for 2 h. The reaction was concentrated and purified by column chromatography (5% methanol/dichloromethane to 7:93:1 methanol/dichloromethane/33% trimethylamine in ethanol) gave 89.6 mg (97%) as a white solid. Mass spec.: 755.37 (MH)<sup>+</sup>.

15 (R)-1-Methoxy-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxopropan-2-yl 4-(8-fluoro-2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate

- 111 -




(R)-Methyl 3-(4-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-6-yl)-2-((4-nitrophenoxy)carbonyloxy)propanoate (75 mg, 0.24 mmol) was dissolved in *N,N*-dimethylformamide (6 mL). *N,N*-Diisopropylethylamine (1.5 mL, 8.6 mmol) was added to the mixture followed by 8-fluoro-3-(piperidin-4-yl)quinolin-2(1H)-one hydrochloride (75 mg, 0.27 mmol). The reaction stirred at room temperature for 3 h. The mixture was diluted with ethyl acetate (20 mL), washed successively with water (2X), 1 N hydrochloric acid, and brine. The organic layer was dried (magnesium sulfate), filtered, and concentrated *in vacuo*. The title compound was obtained as white solid in 82% yield and used without further purification. Mass spec.: 637.2 (MH)<sup>+</sup>.

(R)-2-(4-(8-Fluoro-2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carbonyloxy)-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)propanoic acid




- 112 -



(R)-1-Methoxy-3-(4-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-6-yl)-1-oxopropan-2-yl 4-(8-fluoro-2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate (120 mg, 0.19 mmol) was dissolved in tetrahydrofuran (3 mL). Water (2 mL) was added to the mixture followed by lithium hydroxide monohydrate (25 mg, 0.60 mmol). The reaction was stirred at room temperature for 3 h and quenched by the addition of 1 N hydrochloric acid. The mixture was extracted with ethyl acetate (2X). The combined organics were dried (magnesium sulfate), filtered, and concentrated *in vacuo*. The title compound was obtained without further purification as white solid in 95% yield. Mass spec.: 621.1 (M-H)<sup>+</sup>.

(R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(8-fluoro-2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate

- 113 -




(R)-2-(4-(8-Fluoro-2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxyloxy)-3-(7-methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)propanoic acid (105 mg, 0.17 mmol) was dissolved in *N,N*-dimethylformamide (3 mL). *N,N*-Diisopropylethylamine (100  $\mu$ L, 0.57 mmol) was added to the mixture followed by *o*-benzotriazol-1-yl-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (65 mg, 0.20 mmol). The reaction was stirred at room temperature for 5 min. 4-Piperidinopiperidine (41 mg, 0.22 mmol) was added to the mixture. The reaction was stirred at room temperature for 2 h. The mixture was diluted with ethyl acetate (25 mL), washed successively with water (3X), and brine. The organic layer was dried (magnesium sulfate), filtered, and concentrated *in vacuo*. The title compound was obtained without further purification as white solid in 96% yield. Mass spec.: 773.3 (MH)<sup>+</sup>.

15

### Example 15

(R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(8-fluoro-2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate


- 114 -



(R)-3-(7-Methyl-2-((2-(trimethylsilyl)ethoxy)methyl)-2H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(8-fluoro-2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate (122 mg, 0.16 mmol) was dissolved in ethyl acetate (2 mL). Hydrochloric acid (4 N in dioxane, 3.0 mL) was added to the mixture. The reaction was stirred at room temperature for 18 h. The reaction mixture was concentrated *en vacuo* and purified by preparatory HPLC. Organic solvents were removed from the product fractions. The remaining aqueous solution was made basic with aqueous sodium bicarbonate and was extracted with ethyl acetate (2X). The combined organic extracts were dried (magnesium sulfate), filtered, and concentrated *in vacuo*. The title compound was obtained as an off-white solid in 59% yield. <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>): δ 13.03 (s, 1H), 11.78 (s, 1H), 7.99 (s, 1H), 7.70 (m, 1H), 7.46 (m, 3H), 7.34 (dd, *J*=10.81, 8.09, 1H), 7.15 (m, 1H), 7.06 (s, 1H), 5.43 (m, 1H), 4.34 (m, 1H), 4.17 (m, 1H), 3.88 (m, 1H), 3.06 (m, 2H), 2.97 (m, 2H), 2.83 (m, 3H), 2.50 (m, 3H), 2.36 (m, 2H), 2.27 (m, 1H), 2.05 (m, 2H), 1.82 (d, *J*=12.51, 2H), 1.68 (m, 1H), 1.39 (m, 10H), 0.78-0.26 (m, 1H). Mass spec.: 643.3425 (MH)<sup>+</sup>.


20 *tert*-Butyl 2-fluorophenylcarbamate

- 115 -



To a solution of di-*tert*-butyldicarbonate (90.4 g, 414 mmol) in tetrahydrofuran (414 mL) was added 2-fluorobenzenamine (40.0 mL, 414 mmol). The reaction was heated at reflux overnight. The reaction was cooled, concentrated, dissolved in pentane, washed, in order, with 1 N potassium bisulfate (2X); water, 20% potassium hydroxide, and brine, and then dried over magnesium sulfate and concentrated to give a light brown oil which was dried under high vacuum to give 83.7 g (96%) as a light brown oil which was used without further purification. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 1.52 (s, 9H), 6.68 (bs, 1H), 6.85-7.20 (m, 3H), 8.07 (dd, *J*=8.1, 8.1 Hz, 1H). Mass spec.: 234.18 (MNa)<sup>+</sup>.

*tert*-Butyl 2-fluoro-6-formylphenylcarbamate



15


*tert*-Butyl 2-fluorophenylcarbamate (42.7 g, 202 mmol) was concentrated from toluene *in vacuo* (3X) to remove any traces of water. The resulting residue was dissolved in tetrahydrofuran (600 mL), and cooled to -78°C. To this was added *tert*-butyllithium (1.7 M in pentane, 285 mL, 485 mmol) in dropwise fashion. After addition was complete, the reaction was stirred at -78°C for 30 minutes. The solution was allowed to gradually warm to -20°C before re-cooling to -78°C. To this was added dimethylformamide (46.9 mL, 606 mmol). The reaction was allowed to gradually warm to room temperature overnight. The

- 116 -

reaction was poured into a separatory funnel containing diethyl ether and water. The organics were washed with water (3X) and concentrated. The combined aqueous washes were neutralized with 1 M potassium bisulfate and extracted with diethyl ether. The ethereal layers were combined with the other organics, washed with water, then brine, dried over magnesium sulfate, and concentrated to give 48.1 g (100%) as a yellow oil which was used without further purification. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 1.50 (s, 9H), 7.27 (m, 1H), 7.35 (m, 1H), 7.56 (d, *J*=7.6 Hz, 1H), 7.85 (bs, 1H), 9.99 (d, *J*=1.2 Hz, 1H). Mass spec.: 262.16 (MNa)<sup>+</sup>.

10

*tert*-Butyl 2-((1-benzylpiperidin-4-ylamino)methyl)-6-fluorophenylcarbamate




To a solution of *tert*-butyl 2-fluoro-6-formylphenylcarbamate (48.0 g, 201 mmol) in ethanol (100 mL) was added 4-amino-1-benzylpiperidine (41.0 mL, 201 mmol). The solution was concentrated *in vacuo*, and then water was removed by twice dissolving the residue in toluene and concentrating the solution *in vacuo*. The resulting oil was dissolved in tetrahydrofuran (250 mL), cooled to 0°C, and treated with sodium borohydride (3.80 g, 100 mmol). The ice bath was removed and stirring continued overnight. The reaction was treated with ethanol (250 mL), an additional portion of sodium borohydride (2.00 g, 53 mmol), and an additional portion of 4-amino-1-benzylpiperidine (2.0 mL, 9.8 mmol). The resulting solution was stirred for 4 h at room temperature. The reaction was cooled to 0°C, quenched by addition of saturated ammonium chloride, filtered to remove solids, and concentrated to remove most (but not all) of the tetrahydrofuran. The reaction was extracted with diethyl ether (2X). The ethereal

- 117 -


layer was washed with water (3X), then brine, dried over magnesium sulfate, and concentrated to give 83 g (100%) as a viscous yellow oil which was pure enough to use in the following step. Mass spec.: 414.51 (MH)<sup>+</sup>.

5 3-(1-Benzylpiperidin-4-yl)-8-fluoro-3,4-dihydroquinazolin-2(1H)-one



*tert*-Butyl 2-((1-benzylpiperidin-4-ylamino)methyl)-6-fluorophenylcarbamate (83.0 g, 201 mmol) was dissolved in pyridine (600 mL) and heated at reflux for 12 h. The reaction was concentrated, triturated with hot diethyl ether and placed in the freezer overnight. The resulting solid was filtered to give 68.1 g (64%) as a white solid. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 1.68 (m, 2H), 1.86 (dddd, *J*=11.9, 11.9, 11.9, 3.4 Hz, 2H), 2.14 (dd, *J*=11.6, 10.1 Hz, 2H), 2.98 (d, *J*=11.6 Hz, 2H), 3.51 (s, 2H), 4.34-4.44 (m, 3H), 6.71 (bs, 1H), 6.79-6.89 (m, 2H), 6.94 (dd, *J*=9.2, 9.2 Hz, 1H), 7.21-7.34 (m, 5H). Mass spec.: 340.30 (MH)<sup>+</sup>.

8-Fluoro-3-(piperidin-4-yl)-3,4-dihydroquinazolin-2(1H)-one



20

To a solution of 3-(1-benzylpiperidin-4-yl)-8-fluoro-3,4-dihydroquinazolin-2(1H)-one (2.50 g, 7.37 mmol) in acetic acid (50 mL) under nitrogen, was added palladium (10% on charcoal, 300 mg). The Parr shaker was pressurized to 50 psi and shaken for two days. The flask was flushed with nitrogen, filtered, and concentrated. The residue was dissolved in methanol and


25

- 118 -

re-concentrated to give a light brown oil which solidified upon standing. The residue was dissolved in 1 N hydrochloric acid, and concentrated *in vacuo*, maintaining the temperature at 40°C. The resulting oil was dissolved in methanol, concentrated, and dried under high vacuum to give 2.19 g (quant.) of the hydrochloride salt as an off-white solid.

Generation of the freebase: The hydrochloride salt (15.1 g, 52.8 mmol) was suspended in 2 N sodium hydroxide (40 mL) and stirred at room temperature for 2.5 h. The remaining solid was filtered, washed with water (0°C, 2X 50 mL), then anhydrous diethyl ether (100 mL). The resulting solid was dried under high vacuum overnight to give 12.5 g (95%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz) δ 1.71 (m, 4H), 2.75 (m, 2H), 3.16 (m, 2H), 4.38 (s, 2H), 4.46 (m, 1H), 6.77 (bs, 1H), 6.81-6.89 (m, 2H), 6.95 (m, 1H). Mass spec.: 250.22 (MH)<sup>+</sup>.

Methyl 2-(1-benzylpiperidin-4-yl)acetate




Sodium hydride (60% in mineral oil, 10.55 g, 264 mmol) was washed with hexanes then suspended in *N,N*-dimethylformamide (200 mL). Mixture was cooled to 0°C. Trimethyl phosphonoacetate (38.0 mL, 249 mmol) was added to the mixture dropwise. The reaction was stirred at 0°C for 30 minutes. 1-Benzyl-4-piperidone (40.0 mL, 220 mmol) was added to the reaction mixture dropwise. The reaction was warmed to ambient temperature and held with stirring for 1 h. The reaction mixture was diluted with diethyl ether (500 mL), washed with water (2X), then brine. The organic layer was dried (magnesium sulfate), filtered, and concentrated *in vacuo*. The residue was dissolved in methanol (220 mL). Platinum(IV) oxide (600 mg, 2.64 mmol) was added to the mixture. The reaction vessel was placed on a Parr apparatus, charged with 40 psi of hydrogen gas, and shaken at room temperature for 5 h. The reaction mixture was removed from the apparatus, filtered through celite, and concentrated. The residue was passed through a short column of silica gel eluting with ethyl acetate. Fractions were

- 119 -

concentrated *in vacuo*. The title compound was obtained as amber oil in 90% yield.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.31-7.16 (m, 5H), 3.62 (s, 3H), 3.45 (s, 2H), 2.83 (d,  $J=11.71$ , 2H), 2.20 (d,  $J=6.95$ , 2H), 2.00-1.88 (m, 1H), 1.82-1.69 (m, 1H), 1.69-1.59 (m, 2H), 1.38-1.25 (m, 2H). Mass spec.: 249.3 (MH) $^+$ .


5

Methyl 2-(1-benzylpiperidin-4-yl)-3-hydroxy-3-(2-nitrophenyl)propanoate



Diisopropylamine (3.50 mL, 24.9 mmol) was dissolved in tetrahydrofuran  
10 (30 mL). The mixture was cooled to  $-78^\circ\text{C}$ . Butyllithium (2.5 M in pentane, 9.8 mL, 24.5 mmol) was added to the mixture dropwise, and the reaction stirred at  $-78^\circ\text{C}$  for 15 min. A solution of methyl 2-(1-benzylpiperidin-4-yl)acetate (5.50 g, 22.2 mmol) in THF (8 mL) was then added to the mixture dropwise over 20 minutes. The reaction was stirred at  $-78^\circ\text{C}$  for 45 minutes. A solution of 2-nitrobenzaldehyde (3.70 g, 24.5 mmol) in THF (5 mL) was then added to the  
15 mixture dropwise over 15 minutes. The reaction was stirred at  $-78^\circ\text{C}$  for 30 minutes and quenched by the addition of saturated aqueous ammonium chloride. The resulting mixture was warmed to room temperature, extracted with ethyl acetate (2X). The combined organics were dried (magnesium sulfate), filtered,  
20 and concentrated. Silica gel chromatography afforded the desired product in 89% yield as light yellow foam. Mass spec.: 399.3 (MH) $^+$ .

3-(1-Benzylpiperidin-4-yl)-4-hydroxy-3,4-dihydroquinolin-2(1H)-one






- 120 -

Methyl 2-(1-benzylpiperidin-4-yl)-3-hydroxy-3-(2-nitrophenyl)propanoate (950 mg, 2.4 mmol) was dissolved in acetic acid (20 mL). Iron(0) (1.0 g, 17.7 mmol) was added to the mixture. The reaction was heated at 85°C and held with stirring for 1.5 h. The mixture was cooled to room temperature and diluted with water (30 mL). The liquid was decanted away from the solids. The aqueous solution was concentrated *in vacuo*. The residue was treated with ethyl acetate (50 mL). The mixture was made basic with aqueous sodium hydroxide. Celite was added to the resulting suspension to create a slurry which was in turn was filtered. The filtrate layers were separated. The aqueous layer was extracted with ethyl acetate. Combined organic layers were dried (magnesium sulfate), filtered, and concentrated *in vacuo*. The title compound was obtained without further purification as yellow oil in 69% yield. Mass spec.: 335.3 (MH)<sup>+</sup>.

3-(1-Benzylpiperidin-4-yl)quinolin-2(1H)-one




3-(1-Benzylpiperidin-4-yl)-4-hydroxy-3,4-dihydroquinolin-2(1H)-one (550 mg, 1.6 mmol) was suspended in benzene (10 mL). *p*-Toluenesulfonic acid monohydrate (370 mg, 1.9 mmol) was added to the mixture. The reaction was heated to reflux and held there for 1 h. The reaction mixture was concentrated *in vacuo*. The resulting residue was dissolved in 10% ethanol/dichloromethane (50 mL) and washed with aqueous sodium bicarbonate (2X). The organic layer was dried (magnesium sulfate), filtered, and concentrated *in vacuo*. The residue was triturated with diethyl ether to give a solid which was filtered, washed with diethyl ether, and dried *in vacuo*. The title compound was obtained as off-white solid in 63% yield. <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 11.72 (s, 1H), 7.72 (s, 1H), 7.62 (d, *J*=6.95, 1H), 7.47-7.38 (m, 1H), 7.35-7.30 (m, 4H), 7.29-7.20 (m,

- 121 -


2H), 7.14 (t,  $J=7.50$ , 1H), 3.49 (s, 3H), 2.92 (d,  $J=11.34$ , 2H), 2.83-2.69 (m, 1H), 2.04 (t,  $J=10.61$ , 2H), 1.78 (d,  $J=12.08$ , 2H), 1.71-1.47 (m, 2H). Mass spec.: 319.3 (MH)<sup>+</sup>.

5 3-(Piperidin-4-yl)quinolin-2(1H)-one



3-(1-Benzylpiperidin-4-yl)quinolin-2(1H)-one (1.72 g, 5.40 mmol) was suspended in methanol (70 mL). A catalytic amount of palladium hydroxide (20% on carbon) was added to the mixture. The reaction vessel was placed on a Parr apparatus and charged with 55 psi of hydrogen. The reaction was shaken at room temperature for 5 h. The mixture was removed from the apparatus and filtered. The filtrate was concentrated to give the title compound as white solid in 90% yield. <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 7.65 (s, 1H), 7.64 (d,  $J=10.61$ , 1H), 7.41 (t,  $J=7.50$ , 1H), 7.26 (d,  $J=8.05$ , 1H), 7.13 (t,  $J=7.32$ , 1H), 3.02 (d,  $J=11.71$ , 2H), 2.82 (t,  $J=11.89$ , 2H), 2.58 (t,  $J=11.71$ , 2H), 1.73 (t,  $J=11.71$ , 2H), 1.42 (m, 2H). Mass spec.: 229.4 (MH)<sup>+</sup>.

*N*-(2-Bromo-6-fluorophenyl)pivalamide



20


2-Bromo-6-fluoroaniline (8.2 g, 43.2 mmol) was dissolved in pyridine (10 mL) and treated with pivaloyl chloride (7.0 mL, 57.2 mmol). The reaction was stirred at room temperature for 3 h. The reaction mixture was concentrated *in vacuo* and treated with ethyl acetate (50 mL). Mixture was washed 1 N

25

- 122 -

hydrochloric acid (2X), then brine. The organic layer was dried (magnesium sulfate), filtered, and concentrated *in vacuo*. The residue was triturated with hexanes to give a solid which was filtered, washed with hexanes, and dried *in vacuo*. The title compound was obtained as white solid in 76% yield. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 7.39-7.31 (m, 1H), 7.14-7.03 (m, 2H), 6.98 (bs, 1H), 1.34 (s, 9H). Mass spec.: 274.1 (MH)<sup>+</sup>, 276.1 (MNa)<sup>+</sup>.

*N*-(2-Fluoro-6-formylphenyl)pivalamide




10

*N*-(2-Bromo-6-fluorophenyl)pivalamide (7.0 g, 25.5 mmol) was dissolved in tetrahydrofuran (200 mL). The mixture was cooled to -78°C, and treated with butyllithium (2 M in cyclohexane, 31.0 mL, 62.0 mmol) dropwise. The reaction mixture was held at -78°C for 30 minutes. A solution of *N,N*-dimethylformamide (10.0 mL, 129 mmol) in tetrahydrofuran (30 mL) was added to the reaction mixture dropwise. The reaction was held at -78°C for 30 minutes and quenched by the addition of aqueous ammonium chloride. The mixture was allowed to warm to room temperature and extracted with ethyl acetate (2X). The combined organic layers were dried (magnesium sulfate), filtered, and concentrated. Silica gel chromatography afforded the desired product as white solid in 80% yield. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 9.93 (d, *J*=1.83, 1H), 9.14 (bs, 1H), 7.56-7.50 (m, 1H), 7.42-7.25 (m, 2H), 1.34 (s, 9H). Mass spec.: 224.2 (MH)<sup>+</sup>.

20

*tert*-Butyl 4-(2-methoxy-2-oxoethyl)piperidine-1-carboxylate




25

- 123 -

Sodium hydride (60% in mineral oil, 4.8 g, 120 mmol) was washed with hexanes and then suspended in *N,N*-dimethylformamide (100 mL). The mixture was cooled to 0°C. Trimethyl phosphonoacetate (17.0 mL, 111 mmol) was added to the mixture dropwise. The reaction was held at 0°C for 45 minutes. A solution of *N-tert*-butoxycarbonyl-4-piperidone (18.5 g, 92.6 mmol) in *N,N*-dimethylformamide (25 mL) was added to the reaction mixture dropwise. The mixture was held at 0°C for 1 h and then warmed to room temperature where it was held for 1 h. The reaction was quenched with 1 N hydrochloric acid. The mixture was extracted with diethyl ether (2X). Combined organic layers were washed with water (2X), then brine. The organic layer was dried (magnesium sulfate), filtered, and concentrated *in vacuo*. The residue was dissolved in 1:1 ethyl acetate/methanol (60 mL). A catalytic amount of palladium (10% on charcoal) was added to the mixture. The reaction vessel was placed on a Parr apparatus, charged with 55 psi of hydrogen, and shaken at room temperature for 18 h. The reaction mixture was removed from the Parr apparatus and filtered. The filtrate was concentrated *in vacuo* to give the title compound as lightly colored oil in 94% yield. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 4.04 (d, *J*= 10.25, 2H), 3.64 (s, 3H), 2.68 (t, *J*= 14.44, 2H), 2.21 (d, *J*=6.95, 2H), 1.99-1.80 (m, 1H), 1.64 (d, *J*= 13.54, 2H), 1.41 (s, 9H), 1.25-1.03 (m, 2H).

*tert*-Butyl 4-(1-(3-fluoro-2-pivalamidophenyl)-1-hydroxy-3-methoxy-3-oxopropan-2-yl)piperidine-1-carboxylate




- 124 -

Diisopropylamine (3.40 mL, 24.2 mmol) was dissolved in tetrahydrofuran (70 mL). The mixture was cooled to -78°C. Butyllithium (2 M in cyclohexane, 12.2 mL, 24.4 mmol) was added dropwise to the reaction. The mixture was held at -78°C with stirring and held for 20 minutes. A solution of *tert*-butyl 4-(2-methoxy-2-oxoethyl)piperidine-1-carboxylate (5.20 g, 20.2 mmol) in tetrahydrofuran (15 mL) was added to the mixture dropwise. The mixture was held at -78°C with stirring and held for 45 min. In a separate flask, sodium hydride (60% in mineral oil, 970 mg, 24.3 mmol) was washed with hexanes then suspended in tetrahydrofuran (50 mL). The mixture was cooled to 0°C. A solution of *N*-(2-Fluoro-6-formylphenyl)pivalamide (4.50 g, 20.2 mmol) in tetrahydrofuran (20 mL) was added to the mixture dropwise. The mixture was held at 0°C with stirring and held for 1 h. The above prepared aldehyde mixture was added to the ester mixture dropwise over 1.25 h. The mixture was held at -78°C with stirring and held for 1 h. The reaction was quenched with aqueous ammonium chloride, warmed to room temperature, and diluted with water. The mixture was extracted ethyl acetate (2X) and the aqueous phase was discarded. The material was dried (magnesium sulfate), filtered, and concentrated to dryness. Silica gel chromatography gave the title compound as white foam in 81% yield. Mass spec.: 381.2 ( $M-C_4H_8O_2+H$ )<sup>+</sup>.

20

8-Fluoro-3-(piperidin-4-yl)quinolin-2(1H)-one hydrochloride




*tert*-Butyl 4-(1-(3-fluoro-2-pivalamidophenyl)-1-hydroxy-3-methoxy-3-oxopropan-2-yl)piperidine-1-carboxylate (7.86 g, 16.4 mmol) was dissolved in methanol (20 mL). Water (45 mL) was added to the mixture followed by concentrated hydrochloric acid (15 mL, 183 mmol). The reaction was heated to

25

- 125 -


reflux and held for 2.5 h. The reaction mixture was concentrated *in vacuo*, redissolved in ethanol (50 mL), and concentrated *in vacuo*. The residue was crystallized from ethanol. The resulting solids were filtered, washed with cold ethanol, and dried *in vacuo*. The title compound was obtained as white solid in  
5 83% yield.  $^1\text{H}$  NMR (500 MHz, DMSO- $d_6$ ):  $\delta$  11.85 (s, 1H), 8.98 (m, 1H), 8.85 (m, 1H), 7.75 (s, 1H), 7.54 (d,  $J=7.63$ , 1H), 7.36 (dd,  $J=10.22$ , 8.09, 1H), 7.21-7.11 (m, 1H), 3.41-3.29 (m, 2H), 3.14-2.94 (m, 3H), 2.02 (d,  $J=13.43$ , 2H), 1.88-1.71 (m, 2H). Mass spec.: 247.2 (MH) $^+$ .

10 *N*-(1-Benzylpiperidin-4-yl)-4-fluoro-2-nitrobenzamide

An oven dried flask was charged with 4-fluoro-2-nitrobenzoic acid (7.87 g, 42.5 mmol), 1-hydroxybenzotriazole (6.32 g, 46.8 mmol), 1-(3-  
15 dimethylaminopropyl)-3-ethylcarbodiimide (8.96 g, 46.7 mmol) and ethyl acetate (150 mL), followed by fast dropwise addition of triethylamine (11.8 mL, 84.7 mmol) at room temperature. The resulting white suspension was stirred for 3 h. The reaction was poured into 1:1 water/ethyl acetate (400 mL). After separation, the aqueous layer was extracted with ethyl acetate (200 mL). The combined  
20 organic layer was washed with brine (50 mL), dried over magnesium sulfate, and concentrated *in vacuo* to give a solid upon standing over night. Trituration and washing with ethyl acetate (2 x 6 mL) afforded the title compound as a light yellow solid (9.87 g, 71% yield).  $^1\text{H}$ -NMR (CDCl $_3$ , 400 MHz)  $\delta$  7.74 (dd,  $J=6.4$ , 2.0, 1H), 7.50 (dd,  $J=6.8$ , 4.0, 1H), 7.43-7.22 (m, 6H), 5.71 (d,  $J=6.4$ , 1H), 4.05-  
25 3.93 (m, 1H), 3.50 (s, 2H), 2.85-2.79 (m, 2H), 2.20-2.14 (m, 2H), 2.05-2.02 (m, 2H), 1.58-1.49 (m, 2H). Mass spec.: 358.49 (MH) $^+$ .

*N*-(2-amino-4-fluorobenzyl)-1-benzylpiperidin-4-amine

- 126 -



To a solution of lithium aluminum hydride (4.01 g, 105.7 mmol) in anhydrous 1,4-dioxane (40 mL) at reflux was added N-(1-benzylpiperidin-4-yl)-4-fluoro-2-nitrobenzamide (9.87 g, 30.2 mmol) in anhydrous 1,4-dioxane (125 mL) dropwise under nitrogen over 30 minutes. The resulting mixture was heated at reflux for 3 h under nitrogen. After cooling to room temperature, the reaction was carefully quenched with ice water (10 mL), sodium hydroxide (50% in water, 50 mL), and extracted with diethyl ether (2 x 500 mL). The combined organic layers were washed with brine (100 mL), dried over magnesium sulfate, and concentrated *in vacuo* to afford the title compound (8.64 g, 100% yield) as a yellow oil, which was pure enough for use in the next step. <sup>1</sup>H-NMR (CD<sub>3</sub>OD, 400 MHz) δ 7.52-7.40 (m, 5H), 7.28-7.18 (m, 1H), 6.53 (dd, *J*=11.2, 2.4, 1H), 6.48-6.38 (m, 1H), 4.33 (s, 2H), 4.18 (s, 2H), 3.68-3.46 (m, 3H), 3.25-3.10 (m, 2H), 2.50-2.42 (m, 2H), 2.16-1.93 (m, 2H). Mass spec.: 314.21 (MH<sup>+</sup>).

3-(1-Benzylpiperidin-4-yl)-7-fluoro-3,4-dihydroquinazolin-2(1H)-one




1,1'-Carbonyldiimidazole (6.92 g, 42.7 mmol) was added to a solution of N-(2-amino-4-fluorobenzyl)-1-benzylpiperidin-4-amine (8.64 g, 30.2 mmol) in dry tetrahydrofuran (150 mL) at 0°C in one portion. After 5 min, the mixture was allowed to warm to room temperature and stirred for 3 h. The reaction was partitioned between water/diethyl ether (200 mL/200 mL). After separation, the aqueous solution was extracted with diethyl ether (200 mL). The combined organic solution was washed with brine (100 mL), dried over magnesium sulfate,

- 127 -

and concentrated *in vacuo* to afford a light yellow solid which was triturated with diethyl ether (3 x 10 mL) to give the title compound as a white solid (3.41 g, 36% yield). <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.40-7.18 (m, 5H), 6.92-6.82 (m, 1H), 6.75 (dd, *J*=8.2, 2.6, 1H), 6.66 (dd, *J*=8.8, 4.8, 1H), 4.48-4.34 (m, 1H), 4.03 (s, 2H),  
 5 3.53 (s, 2H), 2.98 (d, *J*=11.2, 2H), 2.37-2.12 (m, 2H), 1.99-1.82 (m, 2H), 1.67 (d, *J*=11.2, 2H); Mass spec.: 340.03 (MH<sup>+</sup>).

7-Fluoro-3-(piperidin-4-yl)-3,4-dihydroquinazolin-2(1H)-one





10

3-(1-Benzylpiperidin-4-yl)-7-fluoro-3,4-dihydroquinazolin-2(1H)-one (1.32 g, 3.89 mmol) was dissolved in ethanol (100 mL) at room temperature. To this was added palladium (10% on charcoal, 130 mg). The resulting mixture was stirred under a hydrogen balloon for 4 d. The reaction was filtered through a pad  
 15 of celite, eluting with ethanol (50 mL), and concentrated *in vacuo* to afford a yellow residue. Upon addition of diethyl ether (20 mL), a solid precipitated from solution. Trituration and filtration with diethyl ether (2 x 5 mL) gave the title compound as a yellow solid (0.842 g, 87% yield). <sup>1</sup>H-NMR (CD<sub>3</sub>OD, 400 MHz) δ 6.94-6.88 (m, 2H), 6.78 (dd, *J*=6.2, 4.0, 1H), 4.42 (s, 2H), 4.41-4.35 (m, 1H),  
 20 3.53-3.47 (m, 2H), 3.21-3.13 (m, 2H), 2.25-2.14 (m, 2H), 1.98-1.92 (m, 2H). Mass spec.: 250.10 (MH<sup>+</sup>).

Furthermore, the following are also prophetically envisaged as compounds of the present invention (represented by structure):



















- 133 -





- 135 -





- 136 -

### CGRP Binding Assay

*Tissue Culture.* SK-N-MC cells were grown at 37 °C in 5% CO<sub>2</sub> as a monolayer in medium consisting of MEM with Earle's salts and L-glutamine (Gibco) supplemented with 10% fetal bovine serum (Gibco).

5        *Cell Pellets.* The cells were rinsed twice with phosphate-buffered saline (155 mM NaCl, 3.3 mM Na<sub>2</sub>HPO<sub>4</sub>, 1.1 mM KH<sub>2</sub>PO<sub>4</sub>, pH 7.4), and incubated for 5-10 min. at 4 °C in hypotonic lysis buffer consisting of 10 mM Tris (pH 7.4) and 5 mM EDTA. The cells were transferred from plates to polypropylene tubes (16 x 100 mm) and homogenized using a polytron. Homogenates were centrifuged at  
10    32,000 x g for 30 min. The pellets were resuspended in cold hypotonic lysis buffer with 0.1% mammalian protease inhibitor cocktail (Sigma) and assayed for protein concentration. The SK-N-MC homogenate was then aliquoted and stored at -80 °C until needed.


*Radioligand Binding Assay.* The compounds of invention were solubilized  
15    and carried through serial dilutions using 100% DMSO. Aliquots from the compound serial dilutions were further diluted 25 fold into assay buffer (50 mM Tris-Cl pH 7.5, 5 mM MgCl<sub>2</sub>, 0.005% Triton X-100) and transferred (volume 50 µl) into 96 well assay plates. [<sup>125</sup>I]-CGRP (Amersham Biosciences) was diluted to 60 pM in assay buffer and a volume of 50 µl was added to each well. SK-N-MC  
20    pellets were thawed, diluted in assay buffer with fresh 0.1% mammalian protease inhibitor cocktail (Sigma), and homogenized again. SK-N-MC homogenate (5 µg/well) was added in a volume of 100 µl. The assay plates were then incubated at room temperature for 2 h. Assays were stopped by addition of excess cold  
25    wash buffer (20 mM Tris-Cl pH 7.5, 0.1% BSA) immediately followed by filtration over glass fiber filters (Whatman GF/B) previously soaked in 0. 5% PEI. Non-specific binding was defined with 1 µM beta-CGRP. Protein bound radioactivity was determined using a gamma or scintillation counter. The IC<sub>50</sub> was defined as the concentration of a compound of invention required to displace 50% of radioligand binding.

30        Human CGRP receptor binding affinities for Examples 1-15 were each less than 1nM.

- 137 -

What is claimed is:

1. A compound according to Formula (I)



5

or a pharmaceutically acceptable salt or solvate thereof  
wherein

V is  $-N(R^1)(R^2)$  or  $OR^4$ ;

$R^4$  is H,  $C_{1-6}$ alkyl,  $C_{1-4}$ haloalkyl or  $(C_{1-4}$ alkylene) $_{0-1}R^4$

10

$R^{4'}$  is  $C_{3-7}$ cycloalkyl, phenyl, adamantyl, quinuclidyl,

azabicyclo[2.2.1]heptyl, azetidiny, tetrahydrofuranyl,

furanyl, dioxolanyl, thienyl, tetrahydrothienyl, pyrrolyl,

pyrrolinyl, pyrrolidinyl, imidazolyl, imidazoliny,

imidazolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl,

15

oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, oxadiazolyl,

thiadiazolyl, triazolyl, pyranyl, pyridyl, pyrimidinyl,

pyrazinyl, pyridazinyl, triazinyl, piperidinyl, piperazinyl,

morpholino, thiomorpholino or dioxolanyl; and

$R^{4'}$  is optionally substituted with 1 or 2 of the same or

20

different substituents selected from the group

consisting of halo, cyano,  $C_{1-4}$ alkyl,  $C_{1-4}$ haloalkyl,

$C_{1-4}$ alkoxy, hydroxy, amino,  $C_{3-7}$ cycloalkyl,

$C_{1-3}$ alkylamino,  $C_{1-3}$ dialkylamino,

$(C_{1-3}$ alkyl) $_{0-2}$ ureido, phenyl and benzyl; and

25

$R^{4'}$  optionally contains 1 or 2 carbonyls wherein the carbon  
atom of said carbonyl is a member of the ring  
structure of  $R^{4'}$ ;

- 138 -

$R^1$  and  $R^2$  are each independently  $L^1$ , wherein  $L^1$  is selected from the group consisting of H,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $-C_{1-6}$ alkylene-amino( $C_{1-3}$ alkyl)<sub>2</sub>,  $C_{3-7}$ cycloalkyl, phenyl, azetidiny, adamantyl, tetrahydrofuranyl, furanyl, dioxolanyl, thienyl, tetrahydrothienyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, imidazolyl, imidazoliny, imidazolidiny, pyrazolyl, pyrazolinyl, pyrazolidiny, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl, pyranyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, piperidinyl, piperazinyl, morpholino, thiomorpholino and dioxolanyl; and

$R^1$  and  $R^2$  are each optionally and independently substituted with 1 or 2 of the same or different substituents selected from the group consisting of halo, cyano,  $C_{1-4}$ alkyl,  $C_{1-4}$ haloalkyl,  $C_{1-4}$ alkoxy, hydroxy, amino,  $C_{3-7}$ cycloalkyl,  $C_{1-3}$ alkylamino,  $C_{1-3}$ dialkylamino, ( $C_{1-3}$ alkyl)<sub>0-2</sub>ureido, phenyl and benzyl;

$R^1$  and  $R^2$  optionally and independently contain 1 or 2 carbonyls wherein the carbon atom of said carbonyl is a member of the heterocycles comprising  $R^1$  and  $R^2$ ;

wherein  $L^1$  is optionally and independently interrupted from the nitrogen to which it is attached by  $L^2$ , wherein  $L^2$  is independently  $C_{1-3}$ alkylene or  $C_{1-3}$ alkylidene; or

$R^1$  and  $R^2$  together with the nitrogen to which they are attached form X,

wherein X is azetidiny, pyrrolyl, pyrrolinyl, pyrrolidinyl, imidazoliny, imidazolidiny, pyrazolinyl,

- 139 -

pyrazolidinyl, azepinyl, diazepinyl, piperazinyl,  
piperidinyl, morpholino or thiomorpholino;  
wherein X is optionally substituted with Y, wherein

5 Y is dioxolanyl, C<sub>1-9</sub>alkyl, C<sub>2-9</sub>alkenyl,  
C<sub>2-9</sub>alkynyl, C<sub>1-4</sub>alkylamino,  
C<sub>1-4</sub>dialkylamino, C<sub>1-4</sub>alkoxy,  
C<sub>3-7</sub>cycloalkyl, phenyl, azetidiny, furanyl,  
thienyl, pyrrolyl, pyrrolinyl, pyrrolidinyl,  
pyrrolidinonyl, imidazolyl, imidazoliny, 10  
imidazolidinyl, imidazolidinonyl, pyrazolyl,  
pyrazoliny, pyrazolidinyl, azepinyl,  
diazepinyl, pyridyl, pyrimidinyl,  
dihydrobenzimidazolonyl, piperazinyl,  
piperidinyl, morpholino, benzothiazolyl,  
15 benzisothiazolyl or thiomorpholino;  
and wherein X and Y are  
optionally interrupted with Z,  
wherein Z is -NHC(O)O-,  
-NHC(O)NH-, NC(O)NH<sub>2</sub>,  
20 -NH-, -C<sub>1-3</sub>alkylene-,  
-C<sub>1-3</sub>alkylene-,  
-C<sub>1-3</sub>alkenylene-NHC(O)O-  
C<sub>1-3</sub>alkylene-; and  
optionally and independently  
25 substituted with 1 or 2 of the  
same or different substituents  
selected from the group  
consisting of halo, C<sub>1-4</sub>alkyl,  
amino, C<sub>1-3</sub>alkylamino,  
30 -C<sub>1-6</sub>alkylene-  
amino(C<sub>1-3</sub>alkyl)<sub>2</sub>,

- 140 -

(C<sub>1-3</sub>alkyl)<sub>0-2</sub>ureido, phenyl  
and benzyl;

X and Y optionally and  
independently contain 1 or 2  
carbonyls wherein the carbon  
atom of said carbonyl is a  
member of the heterocycles  
comprising X and Y;

provided that if X is substituted with Y, and  
if X and Y are not interrupted with  
Z, then

X and Y optionally share one carbon  
atom and together form a  
spirocyclic moiety;

Q is Q' or Q";  
wherein

Q' is (S<sup>y</sup>)<sub>s</sub>R<sup>3</sup>; and

Q" is NH(S<sup>y</sup>)<sub>s</sub>R<sup>3</sup>, NHC(O)(S<sup>y</sup>)<sub>s</sub>R<sup>3</sup>, NHC(O)O(S<sup>y</sup>)<sub>s</sub>R<sup>3</sup>,  
NHC(O)NH(S<sup>y</sup>)<sub>s</sub>R<sup>3</sup>, O(S<sup>y</sup>)<sub>s</sub>R<sup>3</sup>, (S<sup>y</sup>)<sub>s</sub>NHR<sup>3</sup>,  
(S<sup>y</sup>)<sub>s</sub>NHC(O)R<sup>3</sup>, (S<sup>y</sup>)<sub>s</sub>NHC(O)OR<sup>3</sup>,  
(S<sup>y</sup>)<sub>s</sub>NHC(O)NHR<sup>3</sup> or (S<sup>y</sup>)<sub>s</sub>OR<sup>3</sup>;

wherein S<sup>y</sup> is C<sub>1-3</sub>alkylene or C<sub>1-3</sub>alkylidene and s is  
0 or 1;

R<sup>3</sup> is R<sup>3a</sup> or R<sup>3b</sup>  
wherein

R<sup>3a</sup> is

(vi) a heterocycle having two fused rings with 5 to 7  
members in each of said rings, said heterocycle  
containing one to five of the same or different  
heteroatoms selected from the group consisting of  
O, N and S and said heterocycle optionally

- 141 -

containing 1 or 2 carbonyls wherein the carbon atom of said carbonyl is a member of said fused rings;

(vii) a 4 to 6 membered heterocycle containing one to three of the same or different heteroatoms selected from the group consisting of O, N and S, optionally containing 1 to 2 carbonyls, wherein the carbon atom of said carbonyl is a member of said 4 to 6 membered heterocycle;

(viii) C<sub>3-7</sub>cycloalkyl;

(ix) carbazolyl, fluorenyl, phenyl, -O-phenyl, -O-C<sub>1-4</sub>alkylene-phenyl, or naphthyl; or

(x) C<sub>1-8</sub>alkyl, C<sub>2-7</sub>alkenyl, -C(O)R<sup>3'</sup>, CHC(O)O-R<sup>3'</sup>, CH(CH<sub>3</sub>)C(O)O-R<sup>3'</sup>, -C(O)O-R<sup>3'</sup> or C<sub>2-7</sub>alkynyl; and

wherein R<sup>3a</sup> is optionally substituted with 1 to 3 of the same or different substituents selected from the group consisting of benzyl, phenyl, -O-phenyl, -O-C<sub>1-3</sub>alkylenepheryl, -C<sub>1-3</sub>alkylene-OC(O)-phenyl, cyano, amino, nitro, halo, C<sub>1-6</sub>alkyl, C<sub>1-3</sub>mono-bi-tri-haloalkyl, C<sub>1-3</sub>mono-bi-tri-haloalkyloxy, (C<sub>1-3</sub>alkyl)<sub>1-2</sub>amine, -OR<sup>3'</sup>, -C(O)R<sup>3'</sup>, -C(O)O-R<sup>3'</sup>, -O-C(O)R<sup>3'</sup>, -N(R<sup>3'</sup>)<sub>2</sub>, -C(O)N(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)C(O)(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)C(O)N(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)C(O)OR<sup>3'</sup>, -O-C(O)N(R<sup>3'</sup>)<sub>2</sub>, -N(R<sup>3'</sup>)SO<sub>2</sub>R<sup>3'</sup>, -SO<sub>2</sub>N(R<sup>3'</sup>)<sub>2</sub> and -SO<sub>2</sub>R<sup>3'</sup>;

R<sup>3'</sup> is H or -C<sub>1-6</sub>alkyl;

provided that if R<sup>3a</sup> is , -C(O)R<sup>3'</sup>, CHC(O)O-R<sup>3'</sup>, CH(CH<sub>3</sub>)C(O)O-R<sup>3'</sup> or -C(O)O-R<sup>3'</sup>, then said -C(O)R<sup>3'</sup>, CHC(O)O-R<sup>3'</sup>,

- 142 -

$\text{CH}(\text{CH}_3)\text{C}(\text{O})\text{O}-\text{R}^{3'}$  or  $-\text{C}(\text{O})\text{O}-\text{R}^{3'}$  are  
unsubstituted;

$\text{R}^{3b}$  is  $\text{R}^{3a}$  but is not phenyl, 1-naphthyl, 2-naphthyl,  
1,2,3,4-tetrahydro-1-naphthyl, 1H-indol-3-yl, 1-  
5 methyl-1H-indol-3-yl, 1-formyl-1H-indol-3-yl, 1-  
(1,1-dimethylethoxycarbonyl)-1H-indol-3-yl, 4-  
imidazolyl, 1-methyl-4-imidazolyl, 2-thienyl, 3-  
thienyl, thiazolyl, 1H-indazol-3-yl, 1-methyl-1H-  
indazol-3-yl, benzo[b]fur-3-yl, benzo[b]thien-3-yl,  
10 pyridinyl, quinolinyl or isoquinolinyl; optionally  
substituted in the carbon skeleton with mono-, di-  
or trisubstituted by fluorine, chlorine or bromine  
atoms or by branched or unbranched alkyl groups,  
 $\text{C}_{3-8}$ -cycloalkyl groups, phenylalkyl groups,  
15 alkenyl, alkoxy, phenyl, phenylalkoxy,  
trifluoromethyl, alkoxycarbonylalkyl, carboxyalkyl,  
alkoxycarbonyl, carboxy, dialkylaminoalkyl,  
dialkylaminoalkoxy, hydroxy, nitro, amino,  
acetylamino, propionylamino, benzoyl,  
20 benzoylamino, benzoylmethylamino,  
methylsulphonyloxy, aminocarbonyl,  
alkylaminocarbonyl, dialkylaminocarbonyl,  
alkanoyl, cyano, tetrazolyl, phenyl, pyridinyl,  
thiazolyl, furyl, trifluoromethoxy,  
25 trifluoromethylthio, trifluoromethylsulphinyl- or  
trifluoromethylsulphonyl groups;  
wherein said substituents may be the same or  
different and the above-mentioned benzoyl,  
benzoylamino- and benzoylmethylamino  
30 groups may in turn additionally be  
substituted in the phenyl moiety by a

- 143 -

fluorine, chlorine or bromine atom, or by an alkyl, trifluoromethyl, amino or acetyl amino group;

D is O, NCN or NSO<sub>2</sub>C<sub>1-3</sub>alkyl;

5 A is C, N, CH or COH;

m and n are independently 0, 1 or 2;

provided that

if m and n are 0, then A is not N;

if m is 2, then n is not 2; or

10 if n is 2, then m is not 2;

E is N, CH or C;

p is 0 or 1;

if p is 1, then G, J and E together form A<sup>x</sup> or A<sup>y</sup>;

15 A<sup>x</sup> is a fused heterocycle having two fused rings with 5 to 7 members in each of said rings, said heterocycle containing one to four of the same or different heteroatoms selected from the group consisting of O, N and S; and optionally containing 1 or 2 carbonyls wherein the  
20 carbon atom of said carbonyl is a member of said fused heterocycle;

A<sup>y</sup> is a 4 to 6 membered heterocycle containing one to three heteroatoms selected from the group consisting of O, N and S; and  
25 optionally containing 1 to 2 carbonyls, wherein the carbon atom of said carbonyl is a member of said 4 to 6 membered heterocycle;

wherein A<sup>x</sup> and A<sup>y</sup> are optionally substituted with  
30 C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>haloalkyl, cyano, C<sub>3-7</sub>cycloalkyl, phenyl, halophenyl, halo, furanyl, pyrrolyl, pyrrolinyl, pyrrolidinyl,



- 144 -

imidazolyl, imidazoliny, imidazolidinyl,  
 pyrazolyl, pyrazoliny, pyrazolidinyl,  
 pyridyl, pyrimidinyl, piperidinyl,  
 piperazinyl or morpholino; or

5 if p is 0 such that G and J are each attached to A, then A is C, and  
 G, J and A together form a spirocyclic ring system with  
 said rings of said system containing A and wherein G, J  
 and A together are GJA' or GJA";  
 wherein

10 GJA' is A<sup>x</sup> or A<sup>y</sup>; and  
 GJA" is A<sup>x</sup> or A<sup>y</sup>;  
 provided that

A<sup>x</sup> is not a 1,3-diaza-fused  
 heterocycle; and

15 A<sup>y</sup> is not a 1,3-diaza-heterocycle;

and further provided that

if Q is Q", then R<sup>3</sup> is R<sup>3a</sup>; and

if Q is Q', then

R<sup>3</sup> is R<sup>3b</sup>; or

20 R<sup>3</sup> is R<sup>3a</sup>, p is 0 and G, J and A together form GJA".

2. A compound according to claim 1 wherein Q is Q' and R<sup>3</sup> is R<sup>3b</sup>.

3. A compound according to claim 1 wherein Q is Q', R<sup>3</sup> is R<sup>3a</sup> and p is 0  
 25 such that G, J and A together form GJA".

4. A compound according to claim 1 wherein Q is Q' and Q' is (S<sup>y</sup>)<sub>s</sub>R<sup>3</sup> and s  
 is 0.

30 5. A compound according to claim 1 wherein Q is Q' and Q' is (S<sup>y</sup>)<sub>s</sub>R<sup>3</sup>, S<sup>y</sup> is  
 C<sub>1-3</sub>alkylene and s is 1.

- 145 -

6. A compound according to claim 1 wherein Q is Q' and Q' is (Sy)sR<sub>3</sub>, Sy is methylene and s is 1.
7. A compound according to claim 1 wherein V is -N(R<sup>1</sup>)(R<sup>2</sup>) and wherein  
5 R<sup>1</sup> and R<sup>2</sup> together with the nitrogen to which they are attached form X,  
wherein X is piperidinyl, piperazinyl or morpholino;  
wherein X is substituted with Y, wherein Y is  
10 dioxolanyl, phenyl, pyridyl, piperazinyl,  
piperidinyl or C<sub>1-4</sub>alkyl;  
and wherein X and Y optionally share one  
carbon atom and together form a  
spirocyclic moiety.
8. A compound according to claim 1 wherein V is -N(R<sup>1</sup>)(R<sup>2</sup>) and wherein  
15 R<sup>1</sup> and R<sup>2</sup> together with the nitrogen to which they are attached form X,  
wherein X is piperidinyl, piperazinyl or morpholino;  
wherein X is substituted with Y, wherein Y  
20 is dioxolanyl, phenyl, pyridyl,  
piperazinyl, piperidinyl or C<sub>1-4</sub>alkyl.
9. A compound according to claim 1 wherein X and Y are not interrupted  
25 with Z.
10. A compound according to claim 1 wherein R<sub>3</sub> is R<sub>3a</sub> and R<sub>3a</sub> is phenyl,  
hydroxyphenyl, azetidiny, naphthyl, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl,  
dihydroquinolinonyl, hydroquinolinonyl, quinolinyl, dihydroisoquinolinonyl,  
hydroisoquinolinonyl, isoquinolinyl, dihydroquinazolinonyl,  
30 hydroquinazolinonyl, quinazolinyl, dihydroquinoxalinonyl, hydroquinoxalinonyl,  
quinoxalinyl, benzimidazolyl, indazolyl, dihydrobenzimidazolonyl,

- 146 -

hydrobenzimidazolonyl, benzimidazoliny, dihydro-benzthiazolonyl,  
hydrobenzthiazolonyl, benzthiazolyl, dihydrobenzoxazolyl, benzotriazolyl,  
dihydrobenzothiophenonyl, hydrobenzothiophenonyl, benzothieryl,  
dihydrobenzofuranonyl, hydrobenzofuranonyl, benzofuranyl, benzdioxolanyl,  
5 dihydroindolonyl, hydroindolonyl, indolyl, indoliziny, isoindolyl, indoliny,  
indazolyl, pyrazolyl, pyrazoliny, pyrazolidiny, furanyl, thienyl, pyrrolyl,  
pyrroliny, pyrrolidiny, imidazolyl, imidazoliny, imidazolidiny, pyridyl,  
puriny, carbazolyl, pyrimidiny, piperidiny, triazolopyrimidiny,  
tetrahydropyrazolopyridiny, piperaziny or morpholino; optionally substituted as  
10 provided in the first embodiment of the first aspect.

11. A compound according to claim 1 wherein  $R^3$  is  $R^{3b}$  and  $R^{3b}$  is  
dihydrobenzimidazolonyl, hydrobenzimidazolonyl, benzimidazoliny, dihydro-  
benzthiazolonyl, hydrobenzthiazolonyl, benzthiazolyl,  
15 dihydrobenzothiophenonyl, hydrobenzothiophenonyl, dihydrobenzofuranonyl,  
hydrobenzofuranonyl, 1H-indazol-5-yl, benzdioxolanyl, dihydrobenzoxazolyl,  
benzotriazolyl, dihydroindolonyl, hydroindolonyl, indoliziny, isoindolyl,  
indoliny, pyrazolyl, pyrazoliny, pyrazolidiny, furanyl, pyrrolyl, pyrroliny,  
pyrrolidiny, imidazoliny, imidazolidiny, puriny, carbazolyl, pyrimidiny,  
20 piperidiny, piperaziny or morpholino; optionally substituted as provided in the  
first embodiment of the first aspect.

12. A compound according to claim 1 wherein m and n are each 1.

25 13. A compound according to claim 1 wherein D is O.

14. A compound according to claim 1 wherein p is 0 such that G and J are  
each attached to A, then G, J and A together form a spirocyclic ring system with  
said rings of said system containing A and wherein G, J and A together form a  
30 heterocycle selected from the group consisting of imidazolinonyl,  
imidazolidinonyl, dihydroquinolinonyl, dihydroisoquinolinonyl,

dihydroquinazolinonyl, dihydroquinoxalinonyl, dihydrobenzoxazinyl,  
 hydrobenzoxazinyl, dihydrobenzoxazinonyl, dihydrobenzimidazolonyl,  
 dihydrobenzimidazolyl, dihydro-benzthiazolonyl, dihydrobenzthiazolyl,  
 dihydrobenzothiophenonyl, dihydrobenzofuranonyl, dihydroindolonyl, indoliny,  
 5 pyrazoliny, pyrazolidiny, pyrroliny, pyrrolidiny, imidazoliny, imidazolidiny,  
 piperidiny, piperaziny and morpholino; wherein said heterocycle is optionally  
 substituted with C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>haloalkyl, cyano, C<sub>3-7</sub>cycloalkyl,  
 phenyl, halophenyl, furanyl, pyrrolyl, pyrroliny, pyrrolidiny, imidazolyl,  
 imidazoliny, imidazolidiny, pyrazolyl, pyrazoliny, pyrazolidiny, pyridyl,  
 10 pyrimidiny, piperidiny, piperaziny or morpholino.

15. 3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-  
 yl)propan-2-yl 4-(1,2-dihydro-2-oxoquinazolin-3(4H)-yl)piperidine-1-  
 carboxylate; (R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-  
 15 yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinazolin-3(4H)-  
 yl)piperidine-1-carboxylate; (R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-  
 (piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(8-fluoro-2-oxo-1,2-  
 dihydroquinazolin-3(4H)-yl)piperidine-1-carboxylate; (R)-3-(7-Methyl-1H-  
 indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-1,2-  
 20 dihydroquinolin-3-yl)piperidine-1-carboxylate; (R)-3-(7-Methyl-1H-indazol-5-  
 yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-4-phenyl-2,3-  
 dihydroimidazol-1-yl)piperidine-1-carboxylate; (R)-3-(7-Methyl-1H-indazol-5-  
 yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-hydroxy-4-(2-oxo-  
 1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate; (R)-3-(7-Methyl-1H-indazol-  
 25 5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(7-fluoro-2-oxo-  
 1,2-dihydroquinazolin-3(4H)-yl)piperidine-1-carboxylate; (R)-1-(4-(5,6-  
 Dihydropyridin-1(2H)-yl)piperidin-1-yl)-3-(7-methyl-1H-indazol-5-yl)-1-  
 oxopropan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate; (R)-  
 3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(9-methyl-3,9-diaza-spiro[5.5]undecan-3-  
 30 yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate; (R)-  
 3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(piperidin-1-yl)propan-2-yl 4-(2-oxo-1,2-

dihydroquinolin-3-yl)piperidine-1-carboxylate; (R)-1-(4-Cyclohexylpiperazin-1-yl)-3-(7-methyl-1H-indazol-5-yl)-1-oxopropan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate; (R)-1-(4-(4-Fluorophenyl)piperazin-1-yl)-3-(7-methyl-1H-indazol-5-yl)-1-oxopropan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate; (R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(pyridin-4-yl)piperazin-1-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate; ( $\pm$ )-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate; or (R)-3-(7-Methyl-1H-indazol-5-yl)-1-oxo-1-(4-(piperidin-1-yl)piperidin-1-yl)propan-2-yl 4-(8-fluoro-2-oxo-1,2-dihydroquinolin-3-yl)piperidine-1-carboxylate or pharmaceutically acceptable salts or solvates thereof.

16. A pharmaceutical composition comprising a compound according to claim 1.

17. A method of treating migraine comprising administration to a mammal in need thereof an effective anti-migraine amount of a pharmaceutical composition according to claim 16.

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US2005/010330A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C07D401/14 A61K31/44 A61P25/06

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data, BEILSTEIN Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6 552 043 B1 (PATCHETT ARTHUR A ET AL) 22 April 2003 (2003-04-22) column 1, line 9 - line 13; claims 1,5	1-17
X	US 2003/212057 A1 (RUDOLF KLAUS ET AL) 13 November 2003 (2003-11-13) claims 1,8,9	1-17
	----- -/-	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

18 July 2005

Date of mailing of the international search report

29/07/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Seelmann, I

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US2005/010330

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PASTERNAK A ET AL: "Potent, orally bioavailable somatostatin agonists: good absorption achieved by urea backbone cyclization" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, OXFORD, GB, vol. 9, no. 3, 8 February 1999 (1999-02-08), pages 491-496, XP004157253 ISSN: 0960-894X figure 2; compound 4	1
P,X	WO 2004/037810 A (BOEHRINGER INGELHEIM PHARMA GMBH & CO. KG; RUDOLF, KLAUS; MUELLER, STE) 6 May 2004 (2004-05-06) page 89, line 23 - page 94, line 18; claims 1,12,13; examples 13,25,26,30,33,36	1-17

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US2005/010330

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6552043	B1	22-04-2003	AU 6211499 A	17-04-2000
			CA 2345357 A1	06-04-2000
			EP 1117662 A1	25-07-2001
			JP 2002525371 T	13-08-2002
			WO 0018764 A1	06-04-2000
US 2003212057	A1	13-11-2003	DE 19963868 A1	12-07-2001
			AU 3537901 A	16-07-2001
			BG 106848 A	29-12-2002
			BR 0017063 A	22-10-2002
			CA 2395541 A1	12-07-2001
			CN 1414960 A	30-04-2003
			CZ 20022628 A3	16-10-2002
			EA 5600 B1	28-04-2005
			EE 200200369 A	15-10-2003
			WO 0149676 A1	12-07-2001
			EP 1252153 A1	30-10-2002
			HU 0300002 A2	28-05-2003
			JP 2003519222 T	17-06-2003
			MX PA02006509 A	29-11-2002
			NO 20023161 A	28-06-2002
			PL 356400 A1	28-06-2004
			SK 9412002 A3	06-11-2002
			ZA 200205181 A	30-09-2003
WO 2004037810	A	06-05-2004	DE 10250080 A1	13-05-2004
			AU 2003276156 A1	13-05-2004
			WO 2004037810 A1	06-05-2004